

A CENTERLESS VIRASORO ALGEBRA OF MASTER SYMMETRIES FOR THE ABLOWITZ-LADIK HIERARCHY

LUC HAINE AND DIDIER VANDERSTICHELEN

ABSTRACT. We show that the Ablowitz-Ladik hierarchy admits a centerless Virasoro algebra of master symmetries in the sense of Fuchssteiner [16]. An explicit expression for these symmetries is given in terms of a generalization of the Cantero, Moral and Velázquez (CMV) matrices [9] (which can also be used to formulate the hierarchy itself) and their action on the tau-functions of the hierarchy is described. The use of the CMV matrices, in contrast with Hessenberg matrices, turns out to be crucial for obtaining a Lax pair representation of the master symmetries. As an application, a new and transparent derivation of the Virasoro constraints satisfied by the unitary matrix model is presented.

CONTENTS

1. Introduction	2
2. Bi-orthogonal Laurent polynomials and CMV matrices	8
2.1. Bi-orthogonal Laurent polynomials	8
2.2. Five term recurrence relations	11
2.3. Explicit expression for the entries of the CMV matrices	13
3. The AL hierarchy and a Lax pair for its master symmetries	16
3.1. The Ablowitz-Ladik hierarchy	16
3.2. A Lax pair for the master symmetries	20
4. The action of the master symmetries on the tau-functions	25
4.1. Some algebraic lemmas	30
4.2. Expression of the master symmetries on the Plücker coordinates	33
4.3. Action of the Virasoro operators $L_k^{(n)}$ on the Schur polynomials	34
4.4. Proof of the main theorem	42
References	48

Date: August 17 2011.

2010 Mathematics Subject Classification. 37K10, 17B68.

Key words and phrases. Ablowitz-Ladik hierarchy, Master symmetries, Virasoro algebra.

The authors acknowledge the partial support of the Belgian Interuniversity Attraction Pole P06/02.

1. INTRODUCTION

The starting object of our study is the following integral over the group $U(n)$ of unitary $n \times n$ matrices

$$I_n = \int_{U(n)} \det(w(U)) dU,$$

where dU is the standard Haar measure, normalized so that the total volume is 1, and $w(z)$ is some appropriate weight function on the unit circle $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$ which is not necessarily positive or even real valued. The integral appears in combinatorics as well as in random matrix theory for various choices of $w(z)$, see [4, 5, 6, 7, 15, 26, 29] and the references therein. Using the Weyl integral formula, the integral can be rewritten as follows

$$I_n = \frac{1}{n!} \int_{(S^1)^n} |\Delta_n(z)|^2 \prod_{k=1}^n w(z_k) \frac{dz_k}{2\pi i z_k}, \quad (1.1)$$

with $\Delta_n(z)$ the Vandermonde determinant

$$\Delta_n(z) = \prod_{1 \leq k < l \leq n} (z_l - z_k) = \det(z_l^{k-1})_{1 \leq k, l \leq n}.$$

Formula (1.1) for I_n is deeply connected with the theory of bi-orthogonal polynomials on the unit circle, as we briefly recall now. Good references for this material are [4, 5, 15], though these authors don't use the formalism of bi-moment functionals. We denote by $\mathbb{C}[z, z^{-1}]$ the ring of Laurent polynomials over $\mathbb{C}$. A *bi-moment functional* on $\mathbb{C}[z, z^{-1}]$ is a bilinear form

$$\mathcal{L} : \mathbb{C}[z, z^{-1}] \times \mathbb{C}[z, z^{-1}] \rightarrow \mathbb{C}, \quad (f, g) \mapsto \mathcal{L}[f, g]. \quad (1.2)$$

We assume that $\mathcal{L}$ satisfies the *Toeplitz condition*

$$\mathcal{L}[z^n, z^m] = \mathcal{L}[z^{n-m}, 1], \quad \forall n, m \in \mathbb{Z}. \quad (1.3)$$

The example to have in mind in the context of the integral I_n is

$$\mathcal{L}[f, g] = \oint_{S^1} f(z) g(z^{-1}) w(z) \frac{dz}{2\pi i z}.$$

The bi-moments associated to $\mathcal{L}$ are

$$\mu_{mn} = \mathcal{L}[z^m, z^n], \quad \forall m, n \in \mathbb{Z}. \quad (1.4)$$

Because of the Toeplitz condition (1.3), they only depend on the difference $m - n$ and we shall often write

$$\mu_{mn} := \mu_{m-n}. \quad (1.5)$$

In the rest of the paper, we shall freely use both notations for the bi-moments. We shall also assume $\mathcal{L}$ to be *quasi-definite*, that is

$$\det(\mu_{kl})_{0 \leq k, l \leq n-1} \neq 0, \quad \forall n \geq 1. \quad (1.6)$$

This is a necessary and sufficient condition for the existence of a sequence of bi-orthogonal polynomials $\{p_n^{(1)}(z), p_n^{(2)}(z)\}_{n \geq 0}$ with respect to $\mathcal{L}$, that is

$p_n^{(1)}(z)$ and $p_n^{(2)}(z)$ are polynomials of degree n , satisfying the orthogonality conditions

$$\mathcal{L}[p_m^{(1)}(z), p_n^{(2)}(z)] = h_n \delta_{m,n}, \quad h_n \neq 0, \quad \forall m, n \in \mathbb{N}. \quad (1.7)$$

Imposing the polynomials to be monic, one can rewrite the square of the module of the Vandermonde determinant on S^1 as

$$|\Delta_n(z)|^2 = \det(z_l^{k-1})_{1 \leq k, l \leq n} \det(z_l^{1-k})_{1 \leq k, l \leq n}, \quad (1.8)$$

$$= \det(p_{k-1}^{(1)}(z_l))_{1 \leq k, l \leq n} \det(p_{k-1}^{(2)}(z_l^{-1}))_{1 \leq k, l \leq n}. \quad (1.9)$$

By expanding the product of the determinants, using the first equality (1.8), one obtains after some manipulations an expression for (1.1) as a Toeplitz determinant

$$I_n = \det(\mu_{k-l})_{0 \leq k, l \leq n-1}, \quad (1.10)$$

while expanding the product in the second equality (1.9), using the orthogonality relations (1.7), leads to

$$I_n = h_0 h_1 \dots h_{n-1}. \quad (1.11)$$

Introducing the variables

$$x_n = p_n^{(1)}(0), \quad y_n = p_n^{(2)}(0), \quad n \geq 0, \quad (1.12)$$

the monic bi-orthogonal polynomials $\{p_n^{(1)}, p_n^{(2)}\}_{n \geq 0}$ satisfy the Szegő-type coupled linear recurrence relations

$$\begin{aligned} p_{n+1}^{(1)}(z) - zp_n^{(1)}(z) &= x_{n+1} z^n p_n^{(2)}(z^{-1}), \\ p_{n+1}^{(2)}(z) - zp_n^{(2)}(z) &= y_{n+1} z^n p_n^{(1)}(z^{-1}), \end{aligned} \quad (1.13)$$

from which it easily follows that

$$\frac{h_{n+1}}{h_n} = 1 - x_{n+1} y_{n+1}, \quad n \geq 0. \quad (1.14)$$

Combining (1.11) and (1.14) immediately leads to the formula

$$I_n = I_1^n \prod_{i=1}^{n-1} (1 - x_i y_i)^{n-i},$$

which is useful for numerical evaluation of I_n , for instance when x_n, y_n satisfy some discrete type Painlevé equations as studied in [5] and [7].

In this paper, we shall be concerned with the situation when $w(z)$ is a "time dependent" perturbed weight of the following form

$$w(z; t, s) = \rho(z) \exp \left\{ \sum_{j=1}^{\infty} (t_j z^j + s_j z^{-j}) \right\}. \quad (1.15)$$

Then, all the objects introduced above become "time dependent". In [4], Adler and van Moerbeke have shown that the time dependent variables $x_n(t, s), y_n(t, s)$ defined in (1.12) satisfy the (semi-infinite) Ablowitz-Ladik

(AL in short) hierarchy, with $(t, s) = (t_1, t_2, \dots, s_1, s_2, \dots)$ ¹. They also show that the integrals $I_n(t, s)$ are precisely the tau-functions of the hierarchy in the sense of Sato theory and denote them naturally by $\tau_n(t, s)$. As it immediately follows from (1.15), at the level of the bi-moments $\mu_{m,n} = \mu_{m-n}$, the AL hierarchy is given by the simple equations

$$T_k(\mu_j) \equiv \frac{\partial \mu_j}{\partial t_k} = \mu_{j+k}, \quad T_{-k}(\mu_j) \equiv \frac{\partial \mu_j}{\partial s_k} = \mu_{j-k}, \quad \forall k \geq 1. \quad (1.16)$$

Obviously $[T_k, T_l] = 0$, $\forall k, l \in \mathbb{Z}$, if we define $T_0 \mu_j = \mu_j$. Following an idea introduced by Haine and Semengue [20] in the context of the semi-infinite Toda lattice (see also Faybusovich and Gekhtman [14] where the same idea appeared independently), we define the following vector fields on the bi-moments

$$V_k(\mu_j) = (j+k)\mu_{j+k}, \quad \forall k \in \mathbb{Z}. \quad (1.17)$$

These vector fields trivially satisfy the commutation relations

$$[V_k, V_l] = (l-k)V_{k+l}, \quad (1.18)$$

$$[V_k, T_l] = lT_{k+l}, \quad \forall k, l \in \mathbb{Z}, \quad (1.19)$$

from which it follows that

$$[[V_k, T_l], T_l] = l[T_{k+l}, T_l] = 0, \quad \forall k, l \in \mathbb{Z}. \quad (1.20)$$

Equations (1.18), (1.19) and (1.20) mean that the vector fields $V_k, k \in \mathbb{Z}$, form a centerless Virasoro algebra of master symmetries, in the sense of Fuchssteiner [16], for the AL hierarchy. We remind the reader that master symmetries are generators for time dependent symmetries of the hierarchy which are first degree polynomial in the time variables, that is

$$X_{k,l} = V_k + t[V_k, T_l], \quad k \in \mathbb{Z},$$

are time dependent symmetries of the vector field T_l (run with time t) as one immediately checks that

$$\frac{\partial X_{k,l}}{\partial t} + [T_l, X_{k,l}] = [V_k, T_l] + [T_l, V_k + t[V_k, T_l]] = 0,$$

from the commutation relations (1.20). Our main concern in this work will be to translate the action of the master symmetries vector fields defined on the bi-moments by (1.17) to the variables x_n, y_n in which the AL hierarchy is usually formulated, as well as on the tau-functions of the hierarchy.

In [4], the AL hierarchy is described as a reduction of the 2-Toda lattice hierarchy on the pair of Hessenberg matrices

$$l_1 = \begin{pmatrix} -x_1 y_0 & 1 & & & \\ -\frac{h_1}{h_0} x_2 y_0 & -x_2 y_1 & 1 & & O \\ -\frac{h_2}{h_0} x_3 y_0 & -\frac{h_2}{h_1} x_3 y_1 & -x_3 y_2 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

¹In [4, 5] the terminology "Toeplitz hierarchy" instead of "AL hierarchy" is used.

and

$$l_2 = \begin{pmatrix} -x_0y_1 & 1 & & & \\ -\frac{h_1}{h_0}x_0y_2 & -x_1y_2 & 1 & & O \\ -\frac{h_2}{h_0}x_0y_3 & -\frac{h_2}{h_1}x_1y_3 & -x_2y_3 & 1 & \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

which represent the operator of multiplication $\mathbb{C}[z] \rightarrow \mathbb{C}[z] : f(z) \rightarrow zf(z)$ in the bases $p^{(1)}(z) = (p_n^{(1)}(z))_{n \geq 0}$ and $p^{(2)}(z) = (p_n^{(2)}(z))_{n \geq 0}$ of bi-orthogonal polynomials

$$zp^{(1)}(z) = l_1 p^{(1)}(z), \quad zp^{(2)}(z) = l_2 p^{(2)}(z).$$

However, to represent the Virasoro algebra of master symmetries $V_k, k \in \mathbb{Z}$, what we need is a basis of the ring $\mathbb{C}[z, z^{-1}]$ of *Laurent polynomials* in which the operators $d_k = z^{k+1} \frac{d}{dz}, k \in \mathbb{Z}$, satisfying $[d_k, d_l] = (l-k)d_{k+l}$, admit nice matrix representations. Thus, in Section 2, we shall develop a slight generalization of the celebrated Cantero, Moral and Velázquez matrices (CMV matrices in short) introduced in [9]. The generalized CMV matrices are *penta-diagonal* (semi-infinite) matrices A_1, A_2 which will represent multiplication by z in bases of *bi-orthogonal Laurent polynomials*, which will be denoted by $f(z) = (f_n(z))_{n \geq 0}$ and $g(z) = (g_n(z))_{n \geq 0}$, satisfying $\mathcal{L}(f_m, g_n) = \delta_{m,n}h_n$ and the five-term recurrence relations

$$zf(z) = A_1 f(z), \quad zg(z) = A_2 g(z). \quad (1.21)$$

In these bases, we shall have that

$$z^{-1}f(z) = A_1^* f(z), \quad z^{-1}g(z) = A_2^* g(z),$$

with $A_1^* = hA_2^T h^{-1}$, $A_2^* = hA_1^T h^{-1}$ and $h = \text{diag}(h_n)_{n \geq 0}$, so that $A_1^* = A_1^{-1}$ and $A_2^* = A_2^{-1}$. At the end of Section 2, simple expressions for the entries of A_1 and A_2 in terms of $x_n, y_n, n \geq 0$, will be given too. In Section 3, we put this theory to use to obtain Lax pair representations for both the AL hierarchy and its Virasoro algebra of master symmetries. Here we must mention that already Nenciu [25] has obtained a Lax pair representation for the AL hierarchy in terms of CMV matrices, in the special case when $y_n = \overline{x_n}$ (see also Simon [27], [28] for related work). To the best of our knowledge, the use of CMV matrices in the context of the master symmetries, in which they are crucial, is a new result. For a recent account of the huge literature on the AL hierarchy, we refer the reader to Section 3.9 of [17].

In Section 4, we shall translate the action of the master symmetries on the tau-functions. As we already mentioned, the tau functions $\tau_n(t, s)$ of the Ablowitz-Ladik hierarchy are given by the integrals $I_n(t, s)$, which by (1.10) can be expressed as Toeplitz determinants

$$\tau_n(t, s) = \det(\mu_{k-l}(t, s))_{0 \leq k, l < n}. \quad (1.22)$$

The so-called elementary Schur polynomials $S_n(t)$, $t = (t_1, t_2, \dots)$, are defined by the generating function

$$\exp\left(\sum_{k=1}^{\infty} t_k x^k\right) = \sum_{n \in \mathbb{Z}} S_n(t_1, t_2, \dots) x^n. \quad (1.23)$$

By a simple computation, which will be recalled in Section 4, one obtains that the tau functions (1.22) admit the expansion

$$\tau_n(t, s) = \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}} S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s), \quad (1.24)$$

where

$$p_{i_0, \dots, i_{n-1}} = \det(\mu_{i_k - j_l}(0, 0))_{\substack{j_0, \dots, j_{n-1} \\ 0 \leq k, l \leq n-1}}, \quad (1.25)$$

are the so-called Plücker coordinates, and $S_{i_1, \dots, i_k}(t)$ denote the Schur polynomials

$$S_{i_1, \dots, i_k}(t) = \det(S_{i_r + s - r}(t))_{1 \leq r, s \leq k}. \quad (1.26)$$

We will show that the induced action of the master symmetries on the Plücker coordinates of the tau-functions translates into a centerless Virasoro algebra of partial differential operators in the (t, s) variables.

For the convenience of the reader, we summarize below our main results, which will be established respectively in Section 3 and Section 4 of the paper.

Theorem 1.1. *The centerless Virasoro algebra V_k , $k \in \mathbb{Z}$, of master symmetries of the Ablowitz-Ladik hierarchy which are defined on the bi-moments by (1.17), translate as follows on the CMV matrices and the tau-functions of the hierarchy:*

1) *On the CMV matrices (A_1, A_2) , they admit the Lax pair representation*

$$V_k(A_1) = \left[A_1, (D_1 A_1^{k+1})_{--} + (A_1^{k+1} D_1^*)_{--} + k(A_1^k)_{--} \right], \quad \forall k \in \mathbb{Z}, \quad (1.27)$$

$$V_k(A_2) = \left[(D_2 A_2^{1-k})_{--} + (A_2^{1-k} D_2^*)_{--} - k(A_2^{-k})_{--}, A_2 \right], \quad \forall k \in \mathbb{Z}, \quad (1.28)$$

where A_{--} denotes the strictly lower triangular part of A , and D_1 and $(D_1^*)^T$ (respectively D_2 and $(D_2^*)^T$) represent the operator of derivation d/dz in the bases $(f_n(z))_{n \geq 0}$ and $(h_n^{-1} g_n(z^{-1}))_{n \geq 0}$ (respectively $(g_n(z))_{n \geq 0}$ and $(h_n^{-1} f_n(z^{-1}))_{n \geq 0}$), with $f_n(z), g_n(z)$ the bi-orthogonal Laurent polynomials satisfying (1.21) and $\mathcal{L}(f_m, g_n) = h_n \delta_{m,n}$.

2) *On the tau-functions $\tau_n(t, s)$, they are given by a centerless Virasoro algebra of partial differential operators in the (t, s) variables*

$$V_k \tau_n(t, s) = L_k^{(n)} \tau_n(t, s), \quad \forall k \in \mathbb{Z},$$

with $L_k^{(n)}$ defined by

$$L_k^{(n)} = \sum_{j=1}^{k-1} \frac{\partial^2}{\partial t_j \partial t_{k-j}} + n \frac{\partial}{\partial t_k} + \sum_{j=1}^{\infty} j t_j \frac{\partial}{\partial t_{j+k}} - \sum_{j=k+1}^{\infty} j s_j \frac{\partial}{\partial s_{j-k}} - \sum_{j=1}^{k-1} j s_j \frac{\partial}{\partial t_{k-j}} - n k s_k, \quad k \geq 1, \quad (1.29)$$

$$L_0^{(n)} = \sum_{j=1}^{\infty} j t_j \frac{\partial}{\partial t_j} - \sum_{j=1}^{\infty} j s_j \frac{\partial}{\partial s_j}, \quad (1.30)$$

$$L_{-k}^{(n)} = - \sum_{j=1}^{k-1} \frac{\partial^2}{\partial s_j \partial s_{k-j}} - n \frac{\partial}{\partial s_k} - \sum_{j=1}^{\infty} j s_j \frac{\partial}{\partial s_{j+k}} + \sum_{j=k+1}^{\infty} j t_j \frac{\partial}{\partial t_{j-k}} + \sum_{j=1}^{k-1} j t_j \frac{\partial}{\partial s_{k-j}} + n k t_k, \quad k \geq 1, \quad (1.31)$$

satisfying the commutation relations

$$[L_k^{(n)}, L_l^{(n)}] = (k-l)L_{k+l}^{(n)}, \quad \forall k, l \in \mathbb{Z}. \quad (1.32)$$

We like to end with a few comments. First, the general "philosophy" that both the hierarchy and its master symmetries are completely specified by their definition on the bi-moments, rests on a Favard like theorem which states that there is a one-to-one correspondence between CMV matrices (A_1, A_2) , with entries built in terms of x_n and y_n satisfying $x_0 = y_0 = 1$ and $x_n y_n \neq 1, n \geq 1$, and quasi-definite Toeplitz bi-moment functionals defined up to a multiplicative nonzero constant. This theorem can be proven as a generalization to bi-orthogonal Laurent polynomials of a similar result in [10], for orthogonal Laurent polynomials on the unit circle. For a complete and independent proof, see [30]. Second, we feel that there are problems that still need to be investigated in connection with the master symmetries of the AL hierarchy. It is known that there is a strong relation between master symmetries and multi-hamiltonian structures of integrable hierarchies, see [3, 11, 12, 14] and the references therein. However the known results in this direction don't apply in the context of the AL hierarchy. It seems to be related to the existence of a *full* Virasoro algebra of master symmetries, instead of *half* a Virasoro algebra like for the usual Toda lattice hierarchy or the Korteweg-de Vries (KdV) hierarchy. Also as shown by Magri and Zubelli [23], master symmetries of the KdV hierarchy are strongly connected with the Duistermaat-Grünbaum [13] bispectral problem. For similar results in the context of orthogonal polynomials on the line and the Toda lattice, we refer the reader to [18, 19]. Our results prompt towards a study of the bispectral problem in the context of (bi)-orthogonal polynomials on the unit circle, about which little seems to be known.

2. BI-ORTHOGONAL LAURENT POLYNOMIALS AND CMV MATRICES

In [9], Cantero, Moral and Velázquez have constructed two bases of orthogonal Laurent polynomials (in short L-polynomials) for which multiplication by z (and z^{-1}) in the ring $\mathbb{C}[z, z^{-1}]$ is represented by a five-diagonal matrix. When there is an underlying hermitian scalar product satisfying the Toeplitz condition, the two bases which they call right and left orthogonal L-polynomials on S^1 , are obtained by applying the Gram-Schmidt procedure to the ordered bases $\{1, z, z^{-1}, z^2, z^{-2}, \dots\}$ and $\{1, z^{-1}, z, z^{-2}, z^2, \dots\}$ respectively. In [9] the result is obtained for a sesquilinear hermitian quasi-definite form on $\mathbb{C}[z, z^{-1}]$, satisfying the Toeplitz condition. In this section, we show that dropping the condition "hermitian" leads to two sequences of right and left *bi-orthogonal* L-polynomials with the same property. Moreover, we give explicit expressions for the entries of the generalized CMV matrices in terms of the variables $x_n, y_n, n \geq 0$, defined in (1.12).

As with bi-orthogonal polynomials, in all the paper, we shall assume as in (1.2) that $\mathcal{L} : \mathbb{C}[z, z^{-1}] \times \mathbb{C}[z, z^{-1}] \rightarrow \mathbb{C}[z, z^{-1}]$ is a bi-moment functional (i.e. it is bilinear) satisfying the Toeplitz condition (1.3). The case of a sesquilinear form $\langle \cdot | \cdot \rangle$ satisfying the Toeplitz condition can be handled by considering the bi-moment functional $\mathcal{L}[f(z), g(z)] = \langle f(z) | \bar{g}(z) \rangle$, with $\bar{g}(z)$ the Laurent polynomial obtained by conjugating the coefficients of $g(z)$. The two sequences of monic right and left bi-orthogonal L-polynomials we shall construct will be expressed in terms of the sequence of monic bi-orthogonal polynomials $\{p_n^{(1)}(z), p_n^{(2)}(z)\}_{n \geq 0}$, given by the well known formulae

$$p_n^{(1)}(z) = \frac{1}{\tau_n} \det \begin{pmatrix} \mu_{0,0} & \dots & \mu_{0,n-1} & 1 \\ \mu_{1,0} & \dots & \mu_{1,n-1} & z \\ \vdots & & \vdots & \vdots \\ \mu_{n,0} & \dots & \mu_{n,n-1} & z^n \end{pmatrix},$$

$$p_n^{(2)}(z) = \frac{1}{\tau_n} \det \begin{pmatrix} \mu_{0,0} & \mu_{0,1} & \dots & \mu_{0,n} \\ \vdots & \vdots & & \vdots \\ \mu_{n-1,0} & \mu_{n-1,1} & \dots & \mu_{n-1,n} \\ 1 & z & \dots & z^n \end{pmatrix},$$

with $\tau_n = \det (\mu_{kl})_{0 \leq k, l \leq n-1}$.

2.1. Bi-orthogonal Laurent polynomials. The following definition is natural from our previous discussion. We define the vector subspaces

$$\mathbb{L}_{m,n} := \langle z^m, z^{m+1}, \dots, z^{n-1}, z^n \rangle, \quad \forall m, n \in \mathbb{Z}, \quad m \leq n,$$

and for $n \geq 0$

$$\begin{aligned} \mathbb{L}_{2n}^+ &:= \mathbb{L}_{-n,n}, & \mathbb{L}_{2n+1}^+ &:= \mathbb{L}_{-n,n+1}, \\ \mathbb{L}_{2n}^- &:= \mathbb{L}_{-n,n}, & \mathbb{L}_{2n+1}^- &:= \mathbb{L}_{-n-1,n}, \end{aligned}$$

with the convention $\mathbb{L}_{-1}^+ = \mathbb{L}_{-1}^- = \{0\}$.

Definition 2.1. A sequence $\{f_n, g_n\}_{n \geq 0}$ in $\mathbb{C}[z, z^{-1}]$ is a sequence of right (left) bi-orthogonal L-polynomials with respect to $\mathcal{L}$ if

- (1) $f_n, g_n \in \mathbb{L}_n^{+(-)} \setminus \mathbb{L}_{n-1}^{+(-)}$;
- (2) $\mathcal{L}[f_n, g_m] = h_n \delta_{n,m}$, with $h_n \neq 0$.

Remark 2.2. Similarly to orthogonal polynomials, condition (2) in Definition 2.1 can be replaced equivalently by

$$(3r) \quad \begin{cases} \mathcal{L}[f_{2n}, z^k] = 0, \mathcal{L}[z^k, g_{2n}] = 0 & \text{if } -n+1 \leq k \leq n, \\ \mathcal{L}[f_{2n}, z^{-n}] \neq 0, \mathcal{L}[z^{-n}, g_{2n}] \neq 0, \\ \mathcal{L}[f_{2n+1}, z^k] = 0, \mathcal{L}[z^k, g_{2n+1}] = 0 & \text{if } -n \leq k \leq n, \\ \mathcal{L}[f_{2n+1}, z^{n+1}] \neq 0, \mathcal{L}[z^{n+1}, g_{2n+1}] \neq 0, \end{cases}$$

in the case of right bi-orthogonal L-polynomials. For left bi-orthogonal L-polynomials the equivalent condition is

$$(3l) \quad \begin{cases} \mathcal{L}[f_{2n}, z^k] = 0, \mathcal{L}[z^k, g_{2n}] = 0 & \text{if } -n \leq k \leq n-1, \\ \mathcal{L}[f_{2n}, z^n] \neq 0, \mathcal{L}[z^n, g_{2n}] \neq 0, \\ \mathcal{L}[f_{2n+1}, z^k] = 0, \mathcal{L}[z^k, g_{2n+1}] = 0 & \text{if } -n \leq k \leq n, \\ \mathcal{L}[f_{2n+1}, z^{-n-1}] \neq 0, \mathcal{L}[z^{-n-1}, g_{2n+1}] \neq 0. \end{cases}$$

We start by proving that sequences of right and left bi-orthogonal L-polynomials for a given Toeplitz bi-moment functional $\mathcal{L}$ are closely related to each other.

Proposition 2.3. Let $f_n^*(z) = f_n(z^{-1})$ and $g_n^*(z) = g_n(z^{-1})$. Then $\{f_n, g_n\}_{n \geq 0}$ is a sequence of right bi-orthogonal L-polynomials with respect to $\mathcal{L}$ if and only if $\{g_n^*, f_n^*\}_{n \geq 0}$ is a sequence of left bi-orthogonal L-polynomials with respect to $\mathcal{L}$.

Proof. We have $f_n^*, g_n^* \in \mathbb{L}_n^- \setminus \mathbb{L}_{n-1}^-$ if and only if $f_n, g_n \in \mathbb{L}_n^+ \setminus \mathbb{L}_{n-1}^+$. The result then follows from

$$\mathcal{L}[g_m^*(z), f_n^*(z)] = \mathcal{L}[g_m(z^{-1}), f_n(z^{-1})] = \mathcal{L}[f_n(z), g_m(z)].$$

□

Sequences of right or left bi-orthogonal L-polynomials with respect to $\mathcal{L}$ are also very closely related to sequences of bi-orthogonal polynomials for $\mathcal{L}$. This is proven in the next theorem.

Theorem 2.4. Let $\mathcal{L}$ be a Toeplitz bi-moment functional and let $\{f_n, g_n\}_{n \geq 0}$ be a sequence in $\mathbb{C}[z, z^{-1}]$. Let us define

$$\begin{cases} p_{2n}^{(1)}(z) = z^n g_{2n}(z^{-1}), & p_{2n+1}^{(1)}(z) = z^n f_{2n+1}(z), \\ p_{2n}^{(2)}(z) = z^n f_{2n}(z^{-1}), & p_{2n+1}^{(2)}(z) = z^n g_{2n+1}(z). \end{cases} \quad (2.1)$$

The sequence $\{f_n, g_n\}_{n \geq 0}$ is a sequence of right bi-orthogonal L-polynomials with respect to $\mathcal{L}$ if and only if $\{p_n^{(1)}, p_n^{(2)}\}_{n \geq 0}$ is a sequence of bi-orthogonal polynomials with respect to $\mathcal{L}$. Furthermore we have

$$\mathcal{L}[f_n, g_n] = \mathcal{L}[p_n^{(1)}, p_n^{(2)}]. \quad (2.2)$$

An analogous statement holds for sequences $\{f_n, g_n\}_{n \geq 0}$ of left bi-orthogonal L-polynomials, if we define

$$\begin{cases} \tilde{p}_{2n}^{(1)}(z) = z^n f_{2n}(z), & \tilde{p}_{2n+1}^{(1)}(z) = z^n g_{2n+1}(z^{-1}), \\ \tilde{p}_{2n}^{(2)}(z) = z^n g_{2n}(z), & \tilde{p}_{2n+1}^{(2)}(z) = z^n f_{2n+1}(z^{-1}). \end{cases} \quad (2.3)$$

Proof. For $n \geq 0$, we define $\mathbb{P}_n = \langle 1, z, \dots, z^n \rangle$ the vector subspace of polynomials with degree less than or equal to n , and $\mathbb{P}_{-1} := \{0\}$. For $\{p_n^{(1)}, p_n^{(2)}\}_{n \geq 0}$ defined as in (2.1) it is trivial that

$$\begin{aligned} p_{2n}^{(1)}, p_{2n}^{(2)} \in \mathbb{P}_{2n} \setminus \mathbb{P}_{2n-1} &\Leftrightarrow g_{2n}, f_{2n} \in \mathbb{L}_{2n}^+ \setminus \mathbb{L}_{2n-1}^+, \\ p_{2n+1}^{(1)}, p_{2n+1}^{(2)} \in \mathbb{P}_{2n+1} \setminus \mathbb{P}_{2n} &\Leftrightarrow f_{2n+1}, g_{2n+1} \in \mathbb{L}_{2n+1}^+ \setminus \mathbb{L}_{2n}^+. \end{aligned}$$

Furthermore we have using the Toeplitz condition (1.3)

$$\mathcal{L}[p_{2n+1}^{(1)}(z), z^k] = \mathcal{L}[z^n f_{2n+1}(z), z^k] = \mathcal{L}[f_{2n+1}(z), z^{k-n}],$$

and similarly

$$\begin{aligned} \mathcal{L}[p_{2n}^{(1)}(z), z^k] &= \mathcal{L}[z^{n-k}, g_{2n}(z)], & \mathcal{L}[z^k, p_{2n+1}^{(2)}(z)] &= \mathcal{L}[z^{k-n}, g_{2n+1}(z)], \\ \mathcal{L}[z^k, p_{2n}^{(2)}(z)] &= \mathcal{L}[f_{2n}(z), z^{n-k}]. \end{aligned}$$

Consequently we have

$$\begin{aligned} \mathcal{L}[p_{2n+1}^{(1)}(z), z^k] &= 0, \quad 0 \leq k \leq 2n &\Leftrightarrow \mathcal{L}[f_{2n+1}(z), z^k] &= 0, \quad -n \leq k \leq n, \\ \mathcal{L}[p_{2n}^{(1)}(z), z^k] &= 0, \quad 0 \leq k \leq 2n-1 &\Leftrightarrow \mathcal{L}[z^k, g_{2n}(z)] &= 0, \quad -n+1 \leq k \leq n, \\ \mathcal{L}[z^k, p_{2n+1}^{(2)}(z)] &= 0, \quad 0 \leq k \leq 2n &\Leftrightarrow \mathcal{L}[z^k, g_{2n+1}(z)] &= 0, \quad -n \leq k \leq n, \\ \mathcal{L}[z^k, p_{2n}^{(2)}(z)] &= 0, \quad 0 \leq k \leq 2n-1 &\Leftrightarrow \mathcal{L}[f_{2n}(z), z^k] &= 0, \quad -n+1 \leq k \leq n, \end{aligned}$$

and

$$\begin{aligned} \mathcal{L}[p_{2n+1}^{(1)}(z), z^{2n+1}] &\neq 0 &\Leftrightarrow \mathcal{L}[f_{2n+1}(z), z^{n+1}] &\neq 0, \\ \mathcal{L}[p_{2n}^{(1)}(z), z^{2n}] &\neq 0 &\Leftrightarrow \mathcal{L}[z^{-n}, g_{2n}(z)] &\neq 0, \\ \mathcal{L}[z^{2n+1}, p_{2n+1}^{(2)}(z)] &\neq 0 &\Leftrightarrow \mathcal{L}[z^{n+1}, g_{2n+1}(z)] &\neq 0, \\ \mathcal{L}[z^{2n}, p_{2n}^{(2)}(z)] &\neq 0 &\Leftrightarrow \mathcal{L}[f_{2n}(z), z^{-n}] &\neq 0. \end{aligned}$$

Thus, according to Remark 2.2, $\{f_n, g_n\}_{n \geq 0}$ is a sequence of right bi-orthogonal L-polynomials with respect to $\mathcal{L}$ if and only if $\{p_n^{(1)}, p_n^{(2)}\}_{n \geq 0}$ is a sequence of bi-orthogonal polynomials with respect to $\mathcal{L}$. Equation (2.2) follows immediately from the definition (2.1) and the Toeplitz condition (1.3).

The statement (2.3) for sequences of left bi-orthogonal L-polynomials is an immediate consequence of the result for sequences of right bi-orthogonal L-polynomials and Proposition 2.3. This concludes the proof. $\square$

We are now able to prove the existence and the unicity of bi-orthogonal L-polynomials with respect to $\mathcal{L}$.

Corollary 2.5. *Consider a Toeplitz bi-moment functional $\mathcal{L}$. There exist a sequence of right bi-orthogonal L-polynomials with respect to $\mathcal{L}$ and a sequence of left bi-orthogonal L-polynomials with respect to $\mathcal{L}$ if and only if $\mathcal{L}$ is quasi-definite as defined in (1.6). Each L-polynomial in these sequences is uniquely determined up to an arbitrary non-zero factor.*

Proof. By virtue of Theorem 2.4, the existence of a sequence of right or left bi-orthogonal L-polynomials with respect to $\mathcal{L}$ is equivalent to the existence of a sequence of bi-orthogonal polynomials with respect to $\mathcal{L}$, which are known to exist if and only if $\mathcal{L}$ is quasi-definite. Since bi-orthogonal polynomials are uniquely determined up to an arbitrary non-zero factor, the same holds for right and left bi-orthogonal L-polynomials. $\square$

From now on we shall assume that $\mathcal{L}$ is quasi-definite, and $\{f_n, g_n\}_{n \geq 0}$ is a sequence of monic right bi-orthogonal L-polynomials with respect to $\mathcal{L}$, i.e. the coefficients of z^{-n} in f_{2n}, g_{2n} and z^{n+1} in f_{2n+1}, g_{2n+1} are equal to 1. We denote by $\{p_n^{(1)}, p_n^{(2)}\}_{n \geq 0}$ the associated sequence of monic bi-orthogonal polynomials with respect to $\mathcal{L}$, as defined by (2.1).

2.2. Five term recurrence relations. We now prove that bi-orthogonal L-polynomials with respect to a quasi-definite Toeplitz bi-moment functional always satisfy five term recurrence relations. This generalizes the result obtained in [9] for orthogonal L-polynomials associated with a quasi-definite Toeplitz sesquilinear hermitian form. The essential ingredient in the proof in [9] is the Toeplitz condition. Consequently, it can immediately be translated to the case of bi-orthogonal L-polynomials.

Theorem 2.6. *Let $\{f_n, g_n\}_{n \geq 0}$ be a sequence of monic right bi-orthogonal L-polynomials with respect to $\mathcal{L}$, and $f_n^*(z) = f_n(z^{-1})$, $g_n^*(z) = g_n(z^{-1})$. Then for $n \geq 0$ there exist five term recurrence relations*

$$\begin{aligned} z f_n(z) &= \sum_{i=n-2}^{n+2} \alpha_{n,i} f_i(z), & z g_n(z) &= \sum_{i=n-2}^{n+2} \beta_{n,i} g_i(z), \\ z f_n^*(z) &= \sum_{i=n-2}^{n+2} \alpha_{n,i}^* f_i^*(z), & z g_n^*(z) &= \sum_{i=n-2}^{n+2} \beta_{n,i}^* g_i^*(z), \end{aligned}$$

where we use the convention $f_n(z) = g_n(z) = 0$ if $n < 0$, and

$$\alpha_{n,i}^* = \frac{h_n}{h_i} \beta_{i,n}, \quad \beta_{n,i}^* = \frac{h_n}{h_i} \alpha_{i,n},$$

with $h_n = \mathcal{L}[f_n, g_n]$. Moreover, we have for all $n \geq 0$

$$\begin{aligned} \alpha_{2n-1, 2n-3} &= 0, & \alpha_{2n, 2n+2} &= 0, \\ \beta_{2n-1, 2n-3} &= 0, & \beta_{2n, 2n+2} &= 0. \end{aligned}$$

Proof. As $f_n \in \mathbb{L}_n^+ \setminus \mathbb{L}_{n-1}^+$, we have $zf_n(z) \in \mathbb{L}_{n+2}^+$. This implies that zf_n admits an expansion in terms of $f_0, \dots, f_{n+2}$

$$zf_n(z) = \sum_{i=0}^{n+2} \alpha_{n,i} f_i(z),$$

with $\alpha_{n,i} \in \mathbb{C}$, $0 \leq i \leq n+2$. Consequently, by bi-orthogonality of the sequence $\{f_n, g_n\}_{n \geq 0}$ we have

$$\mathcal{L}[zf_n, g_m] = \sum_{i=0}^{n+2} h_i \alpha_{n,i} \delta_{i,m}.$$

But we also have

$$\mathcal{L}[zf_n, zg_k] = \mathcal{L}[f_n, g_k] = 0, \quad 0 \leq k \leq n-1,$$

and $\langle g_0, \dots, g_{n-3} \rangle \subset \langle zg_0, \dots, zg_{n-1} \rangle$. It follows that

$$\mathcal{L}[zf_n, g_k] = 0, \quad 0 \leq k \leq n-3.$$

Consequently we have $\alpha_{n,i} = 0$ if $i < n-2$, and thus

$$zf_n(z) = \sum_{i=n-2}^{n+2} \alpha_{n,i} f_i(z).$$

We prove that $\alpha_{2n,2n+2} = \alpha_{2n-1,2n-3} = 0$. We first prove that $\alpha_{2n,2n+2} = 0$. Indeed, we have $zf_{2n}(z) \in \langle z^{1-n}, \dots, z^{1+n} \rangle$. Consequently, using condition (3r) in Remark 2.2, we have $\mathcal{L}[zf_{2n}, g_{2n+2}] = 0$ and thus $\alpha_{2n,2n+2} = 0$. We also have $\alpha_{2n-1,2n-3} = 0$. Indeed, we have $\mathcal{L}[zf_{2n-1}, g_{2n-3}] = \mathcal{L}[f_{2n-1}, z^{-1}g_{2n-3}]$, and $z^{-1}g_{2n-3}(z) \in \langle z^{1-n}, \dots, z^{n-2} \rangle$. From condition (3r) in Remark 2.2, it follows that $\mathcal{L}[zf_{2n-1}, g_{2n-3}] = 0$. A similar argument gives $\beta_{2n,2n+2} = \beta_{2n-1,2n-3} = 0$. The proof of the other recurrence relations is similar.

The coefficients in the recurrence relations satisfy

$$\begin{aligned} \alpha_{n,i} &= \frac{\mathcal{L}[zf_n, g_i]}{\mathcal{L}[f_i, g_i]}, & \beta_{n,i} &= \frac{\mathcal{L}[f_i, zg_n]}{\mathcal{L}[f_i, g_i]}, \\ \alpha_{n,i}^* &= \frac{\mathcal{L}[g_i^*, zf_n^*]}{\mathcal{L}[g_i^*, f_i^*]}, & \beta_{n,i}^* &= \frac{\mathcal{L}[zg_n^*, f_i^*]}{\mathcal{L}[g_i^*, f_i^*]}. \end{aligned}$$

It follows from the definition of $\{g_n^*, f_n^*\}_{n \geq 0}$ that

$$\alpha_{n,i}^* = \frac{\mathcal{L}[g_i^*, zf_n^*]}{\mathcal{L}[g_i^*, f_i^*]} = \frac{\mathcal{L}[f_n, zg_i]}{\mathcal{L}[f_i, g_i]} = \frac{\mathcal{L}[f_n, zg_i]}{\mathcal{L}[f_n, g_n]} \frac{\mathcal{L}[f_n, g_n]}{\mathcal{L}[f_i, g_i]} = \beta_{i,n} \frac{h_n}{h_i}.$$

Similarly we have

$$\beta_{n,i}^* = \frac{\mathcal{L}[zg_n^*, f_i^*]}{\mathcal{L}[g_i^*, f_i^*]} = \frac{\mathcal{L}[zf_i, g_n]}{\mathcal{L}[f_i, g_i]} = \frac{\mathcal{L}[zf_i, g_n]}{\mathcal{L}[f_n, g_n]} \frac{\mathcal{L}[f_n, g_n]}{\mathcal{L}[f_i, g_i]} = \alpha_{i,n} \frac{h_n}{h_i}.$$

This concludes the proof. $\square$

Corollary 2.7. *With the same notations as in Theorem 2.6 we have*

$$\begin{aligned} z^{-1}f_n(z) &= \sum_{i=n-2}^{n+2} \alpha_{n,i}^* f_i(z), & z^{-1}g_n(z) &= \sum_{i=n-2}^{n+2} \beta_{n,i}^* g_i(z), \\ z^{-1}f_n^*(z) &= \sum_{i=n-2}^{n+2} \alpha_{n,i} f_i^*(z), & z^{-1}g_n^*(z) &= \sum_{i=n-2}^{n+2} \beta_{n,i} g_i^*(z). \end{aligned}$$

Defining the vectors

$$f(z) = (f_n(z))_{n \geq 0}, \quad g(z) = (g_n(z))_{n \geq 0}, \quad (2.4)$$

$$f^*(z) = f(z^{-1}) = (f_n^*(z))_{n \geq 0}, \quad g^*(z) = g(z^{-1}) = (g_n^*(z))_{n \geq 0}, \quad (2.5)$$

the five term recurrence relations obtained in Theorem 2.6 and Corollary 2.7 can be written in vector form

$$\begin{cases} zf(z) = A_1 f(z), \\ zg(z) = A_2 g(z), \\ z^{-1}f(z) = A_1^* f^*(z), \\ z^{-1}g(z) = A_2^* g^*(z), \end{cases} \quad \begin{cases} zf^*(z) = A_1^* f^*(z), \\ zg^*(z) = A_2^* g^*(z), \\ z^{-1}f^*(z) = A_1 f^*(z), \\ z^{-1}g^*(z) = A_2 g^*(z), \end{cases} \quad (2.6)$$

with

$$A_1 = (\alpha_{i,j})_{i,j \geq 0}, \quad A_2 = (\beta_{i,j})_{i,j \geq 0},$$

where $\alpha_{i,j} = \beta_{i,j} = 0$ if $|i - j| > 2$, and

$$A_1^* = h A_2^T h^{-1}, \quad A_2^* = h A_1^T h^{-1}, \quad (2.7)$$

where $h = \text{diag}(h_n)_{n \geq 0}$. We call the matrices A_1, A_2 the CMV matrices. Clearly, from (2.6), we have

$$A_1^* = A_1^{-1}, \quad A_2^* = A_2^{-1}. \quad (2.8)$$

2.3. Explicit expression for the entries of the CMV matrices. Explicit expressions for the entries of the CMV matrices can be found in terms of the variables x_n, y_n introduced in (1.12) defining the Szegő type coupled linear recurrence relations (1.13).

Theorem 2.8. *The non-zero entries of the CMV matrices A_1 and A_2 are*

$$\begin{aligned} (A_1)_{2n-1, 2n+1} &= 1, & (A_1)_{2n-1, 2n-1} &= -x_{2n} y_{2n-1}, \\ (A_1)_{2n-1, 2n} &= -x_{2n+1}, & (A_1)_{2n-1, 2n-2} &= -x_{2n}(1 - x_{2n-1} y_{2n-1}), \end{aligned}$$

$$\begin{aligned} (A_1)_{2n, 2n+1} &= y_{2n}, & (A_1)_{2n, 2n-1} &= y_{2n-1}(1 - x_{2n} y_{2n}), \\ (A_1)_{2n, 2n} &= -x_{2n+1} y_{2n}, & (A_1)_{2n, 2n-2} &= (1 - x_{2n-1} y_{2n-1})(1 - x_{2n} y_{2n}), \end{aligned}$$

and

$$\begin{aligned} (A_2)_{2n-1, 2n+1} &= 1, & (A_2)_{2n-1, 2n-1} &= -x_{2n-1} y_{2n}, \\ (A_2)_{2n-1, 2n} &= -y_{2n+1}, & (A_2)_{2n-1, 2n-2} &= -y_{2n}(1 - x_{2n-1} y_{2n-1}), \end{aligned}$$

$$\begin{aligned} (A_2)_{2n, 2n+1} &= x_{2n}, & (A_2)_{2n, 2n-1} &= x_{2n-1}(1 - x_{2n} y_{2n}), \\ (A_2)_{2n, 2n} &= -x_{2n} y_{2n+1}, & (A_2)_{2n, 2n-2} &= (1 - x_{2n-1} y_{2n-1})(1 - x_{2n} y_{2n}). \end{aligned}$$

Proof. (1) We have

$$(A_1)_{2n-1,2n+1} = \frac{1}{h_{2n+1}} \mathcal{L} [z f_{2n-1}(z), g_{2n+1}(z)].$$

By virtue of Theorem 2.4 we obtain

$$\begin{aligned} (A_1)_{2n-1,2n+1} &= \frac{1}{h_{2n+1}} \mathcal{L} [z^{2-n} p_{2n-1}^{(1)}(z), z^{-n} p_{2n+1}^{(2)}(z)] \\ &= \frac{1}{h_{2n+1}} \mathcal{L} [z^2 p_{2n-1}^{(1)}(z), p_{2n+1}^{(2)}(z)]. \end{aligned}$$

As $z^2 p_{2n-1}^{(1)}(z)$ is a monic polynomial of degree $2n+1$, using the bi-orthogonality of the polynomials, we have

$$(A_1)_{2n-1,2n+1} = \frac{1}{h_{2n+1}} \mathcal{L} [z^{2n+1}, p_{2n+1}^{(2)}(z)] = 1.$$

(2) We have

$$(A_1)_{2n-1,2n} = \frac{1}{h_{2n}} \mathcal{L} [z f_{2n-1}(z), g_{2n}(z)]$$

By virtue of Theorem 2.4 we obtain

$$\begin{aligned} (A_1)_{2n-1,2n} &= \frac{1}{h_{2n}} \mathcal{L} [z^{2-n} p_{2n-1}^{(1)}(z), z^n p_{2n}^{(1)}(z^{-1})] \\ &= \frac{1}{h_{2n}} \mathcal{L} [z^2 p_{2n-1}^{(1)}(z), z^{2n} p_{2n}^{(1)}(z^{-1})]. \end{aligned}$$

By using twice (1.13) we have

$$z^2 p_{2n-1}^{(1)}(z) = p_{2n+1}^{(1)}(z) - x_{2n+1} z^{2n} p_{2n}^{(2)}(z^{-1}) - x_{2n} z^{2n} p_{2n-1}^{(2)}(z^{-1}),$$

and thus

$$\begin{aligned} (A_1)_{2n-1,2n} &= \frac{1}{h_{2n}} \mathcal{L} [p_{2n+1}^{(1)}(z), z^{2n} p_{2n}^{(1)}(z^{-1})] - \frac{x_{2n+1}}{h_{2n}} \mathcal{L} [p_{2n}^{(2)}(z^{-1}), p_{2n}^{(1)}(z^{-1})] \\ &\quad - \frac{x_{2n}}{h_{2n}} \mathcal{L} [p_{2n-1}^{(2)}(z^{-1}), p_{2n}^{(1)}(z^{-1})]. \end{aligned}$$

As $z^{2n} p_{2n}^{(1)}(z^{-1})$ is a polynomial of degree $2n$, the first term is equal to 0 by bi-orthogonality. The remaining terms give

$$\begin{aligned} (A_1)_{2n-1,2n} &= -\frac{x_{2n+1}}{h_{2n}} \mathcal{L} [p_{2n}^{(1)}(z), p_{2n}^{(2)}(z)] - \frac{x_{2n}}{h_{2n}} \mathcal{L} [p_{2n}^{(1)}(z), p_{2n-1}^{(2)}(z)] \\ &= -x_{2n+1}. \end{aligned}$$

(3) We have

$$(A_1)_{2n-1,2n-1} = \frac{1}{h_{2n-1}} \mathcal{L} [z f_{2n-1}(z), g_{2n-1}(z)].$$

By virtue of Theorem 2.4 we obtain

$$\begin{aligned} (A_1)_{2n-1,2n-1} &= \frac{1}{h_{2n-1}} \mathcal{L} [z^{2-n} p_{2n-1}^{(1)}(z), z^{1-n} p_{2n-1}^{(2)}(z)] \\ &= \frac{1}{h_{2n-1}} \mathcal{L} [z p_{2n-1}^{(1)}(z), p_{2n-1}^{(2)}(z)]. \end{aligned}$$

By using (1.13) we have

$$\begin{aligned} (A_1)_{2n-1,2n-1} &= \frac{1}{h_{2n-1}} \mathcal{L} [p_{2n}^{(1)}(z) - x_{2n} z^{2n-1} p_{2n-1}^{(2)}(z^{-1}), p_{2n-1}^{(2)}(z)] \\ &= -\frac{x_{2n}}{h_{2n-1}} \mathcal{L} [z^{2n-1} p_{2n-1}^{(2)}(z^{-1}), p_{2n-1}^{(2)}(z)] \\ &= -\frac{x_{2n}}{h_{2n-1}} \mathcal{L} [y_{2n-1} z^{2n-1}, p_{2n-1}^{(2)}(z)] \\ &= -x_{2n} y_{2n-1}. \end{aligned}$$

(4) We have

$$(A_1)_{2n-1,2n-2} = \frac{1}{h_{2n-2}} \mathcal{L} [z f_{2n-1}(z), g_{2n-2}(z)].$$

By virtue of Theorem 2.4 we obtain

$$\begin{aligned} (A_1)_{2n-1,2n-2} &= \frac{1}{h_{2n-2}} \mathcal{L} [z^{2-n} p_{2n-1}^{(1)}(z), z^{n-1} p_{2n-2}^{(1)}(z^{-1})] \\ &= \frac{1}{h_{2n-2}} \mathcal{L} [z p_{2n-1}^{(1)}(z), z^{2n-2} p_{2n-2}^{(1)}(z^{-1})]. \end{aligned}$$

Using (1.13) we obtain

$$\begin{aligned} (A_1)_{2n-1,2n-2} &= \frac{1}{h_{2n-2}} \mathcal{L} [p_{2n}^{(1)}(z) - x_{2n} z^{2n-1} p_{2n-1}^{(2)}(z^{-1}), z^{2n-2} p_{2n-2}^{(1)}(z^{-1})] \\ &= \frac{1}{h_{2n-2}} \mathcal{L} [p_{2n}^{(1)}(z), z^{2n-2} p_{2n-2}^{(1)}(z^{-1})] - \frac{x_{2n}}{h_{2n-2}} \mathcal{L} [z p_{2n-1}^{(2)}(z^{-1}), p_{2n-2}^{(1)}(z^{-1})]. \end{aligned}$$

The first term is equal to 0 as $z^{2n-2} p_{2n-2}^{(1)}(z^{-1})$ is a polynomial of degree $2n-2$. Consequently we have

$$\begin{aligned} (A_1)_{2n-1,2n-2} &= -\frac{x_{2n}}{h_{2n-2}} \mathcal{L} [z p_{2n-2}^{(1)}(z), p_{2n-1}^{(2)}(z)] \\ &= -\frac{x_{2n}}{h_{2n-2}} \mathcal{L} [z^{2n-1}, p_{2n-1}^{(2)}(z)] \\ &= -\frac{h_{2n-1}}{h_{2n-2}} x_{2n} \\ &= -(1 - x_{2n-1} y_{2n-1}) x_{2n}. \end{aligned}$$

(5) The other relations are proven in a similar way. This finishes the proof. $\square$

3. THE AL HIERARCHY AND A LAX PAIR FOR ITS MASTER SYMMETRIES

In this section we "dress up" the equations defining the Ablowitz-Ladik hierarchy (1.16) and its master symmetries (1.17) on the bi-moments. This leads to Lax pair representations both for the hierarchy and its master symmetries on the CMV matrices. In all this section we shall denote the time variables $(t, s) = (t_1, t_2, \dots, s_1, s_2, \dots)$ of the AL hierarchy by $(t_k)_{k \in \mathbb{Z}}$, with $t_{-k} = s_k, k \geq 1$, and T_0 defined as in the Introduction (see below (1.16)). It is only in the next section that the notation (t, s) will be more convenient.

3.1. The Ablowitz-Ladik hierarchy. Let

$$\chi(z) = (1, z, z^{-1}, z^2, z^{-2}, \dots)^T, \quad (3.1)$$

and let $\mathcal{L}$ be a quasi-definite bi-moment functional satisfying the Toeplitz condition. We introduce two matrices S_1 and S_2 by writing the vectors $f(z), g(z)$ (2.4) of monic right bi-orthogonal L-polynomials with respect to $\mathcal{L}$ as follows

$$f(z) = S_1 \chi(z), \quad g(z) = h (S_2^T)^{-1} \chi(z), \quad (3.2)$$

with $h = \text{diag}(h_n)_{n \geq 0}$ and $h_n = \mathcal{L}[f_n, g_n]$. With this definition, S_1 is a lower triangular matrix with all diagonal elements equal to 1, and S_2 is an upper triangular matrix such that $h^{-1} S_2$ has all diagonal elements equal to 1.

Associated to $\mathcal{L}$ we also define the semi-infinite bi-moment matrix

$$M = \begin{pmatrix} \mu_{0,0} & \mu_{0,1} & \mu_{0,-1} & \cdots \\ \mu_{1,0} & \mu_{1,1} & \mu_{1,-1} & \cdots \\ \mu_{-1,0} & \mu_{-1,1} & \mu_{-1,-1} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \quad (3.3)$$

with $\mu_{m,n}$ as in (1.4), (1.5). The bi-moment matrix M can be written in terms of the vector $\chi(z)$ in (3.1)

$$M = \left(\mathcal{L}[(\chi(z))_m, (\chi(z))_n] \right)_{0 \leq m, n < \infty}.$$

The existence of a sequence of right bi-orthogonal L-polynomials for $\mathcal{L}$ is equivalent to the existence of a factorisation of the bi-moment matrix M in a product of a lower triangular matrix and an upper triangular matrix with non-zero diagonal elements.

Proposition 3.1. *The bi-moment matrix M factorizes in a product of a lower triangular matrix and an upper triangular matrix*

$$M = S_1^{-1} S_2.$$

Proof. By bi-orthogonality of the sequence $\{f_n, g_n\}_{n \geq 0}$, we have

$$\mathcal{L}[f_m, g_n] = h_m \delta_{m,n}.$$

This can be written in matrix form

$$h = (\mathcal{L}[f_m, g_n])_{0 \leq m, n < \infty}.$$

Using the expressions (3.2) we obtain

$$\begin{aligned} h &= \left(\mathcal{L} \left[(S_1 \chi(z))_m, (h (S_2^T)^{-1} \chi(z))_n \right] \right)_{0 \leq m, n \leq \infty} \\ &= S_1 M S_2^{-1} h. \end{aligned}$$

Consequently we have

$$M = S_1^{-1} S_2,$$

which establishes the result. $\square$

We define the semi-infinite shift matrix Λ by

$$\Lambda \chi(z) = z \chi(z). \quad (3.4)$$

We have

$$\Lambda = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 1 & 0 & 0 & \dots \\ 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 1 & \dots \\ 0 & 0 & 1 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}, \quad (3.5)$$

and $\Lambda^{-1} = \Lambda^T$. We leave to the reader to check that, because of the Toeplitz property satisfied by the bi-moments in (3.3), we have the commutation relation

$$[\Lambda, M] = 0. \quad (3.6)$$

The CMV matrices can be obtained by "dressing up" the shift Λ .

Proposition 3.2. *We have*

$$A_1 = S_1 \Lambda S_1^{-1}, \quad A_2 = h (S_2^T)^{-1} \Lambda S_2^T h^{-1}, \quad (3.7)$$

$$A_1^{-1} = S_2 \Lambda^T S_2^{-1}, \quad A_2^{-1} = h (S_1^{-1})^T \Lambda^T S_1^T h^{-1}, \quad (3.8)$$

with S_1 and S_2 defined in (3.2).

Proof. We have

$$A_1 f(z) = z f(z) = z S_1 \chi(z) = S_1 \Lambda \chi(z) = S_1 \Lambda S_1^{-1} f(z).$$

It follows that

$$A_1 = S_1 \Lambda S_1^{-1}.$$

The proof for A_2 is similar. The factorisations in (3.8) follow from (2.7), (2.8) and (3.7). $\square$

Remember that because $\mathcal{L} : \mathbb{C}[z, z^{-1}] \times \mathbb{C}[z, z^{-1}] \rightarrow \mathbb{C}$ is a Toeplitz bi-moment functional, the bi-moments $\mu_{m,n} = \mathcal{L}[z^m, z^n]$ only depend on the difference $m - n$ and can be written as in (1.5) $\mu_{m,n} := \mu_{m-n}$.

The Ablowitz-Ladik hierarchy is defined on the space of bi-moments by the vector fields

$$\boxed{T_k \mu_j \equiv \frac{\partial \mu_j}{\partial t_k} = \mu_{j+k}, \quad \forall k \in \mathbb{Z},} \quad (3.9)$$

where we have put $s_k = t_{-k}$ in (1.16). Obviously, these vector fields satisfy the commutation relations

$$[T_k, T_l] = 0, \quad \forall k, l \in \mathbb{Z}.$$

It follows from the definition of Λ in (3.4) and (3.9) that the time evolution of the bi-moment matrix M is given by the equations

$$\boxed{\frac{\partial M}{\partial t_k} = \Lambda^k M, \quad \forall k \in \mathbb{Z}.} \quad (3.10)$$

Equations (3.9) and (3.10) are two equivalent formulations of the Ablowitz-Ladik vector fields at the level of the bi-moments.

For a square matrix A , we define

- A_0 the diagonal part of A ;
- A_- (resp. A_+) the lower (resp. upper) triangular part of A ;
- A_{--} (resp. A_{++}) the strictly lower (resp. strictly upper) triangular part of A .

We establish the following lemma, based on the factorisation of the moment matrix M in Proposition 3.1 in a product of a lower triangular and an upper triangular matrix.

Lemma 3.3. *We have for $k \in \mathbb{Z}$*

$$\frac{\partial S_1}{\partial t_k} S_1^{-1} = -(A_1^k)_{--}, \quad (3.11)$$

$$(S_2^T h^{-1})^{-1} \frac{\partial (S_2^T h^{-1})}{\partial t_k} = (A_2^{-k})_{--}. \quad (3.12)$$

Proof. On the one hand, we have using Proposition 3.1

$$\frac{\partial M}{\partial t_k} = -S_1^{-1} \frac{\partial S_1}{\partial t_k} S_1^{-1} S_2 + S_1^{-1} \frac{\partial S_2}{\partial t_k}.$$

On the other hand, from equation (3.10) we have

$$\frac{\partial M}{\partial t_k} = \Lambda^k M = \Lambda^k S_1^{-1} S_2.$$

As $A_1 = S_1 \Lambda S_1^{-1}$, we obtain

$$A_1^k = -\frac{\partial S_1}{\partial t_k} S_1^{-1} + \frac{\partial S_2}{\partial t_k} S_2^{-1}.$$

Since $\frac{\partial S_1}{\partial t_k}$ is strictly lower triangular, the first term in the right hand side of this equation is strictly lower triangular. The second term is upper triangular. Consequently, taking the strictly lower triangular part of both sides of the equation yields

$$\frac{\partial S_1}{\partial t_k} S_1^{-1} = -(A_1^k)_{--},$$

which establishes (3.11).

To establish the other formula, we write $M = (S_1^{-1} h) (h^{-1} S_2)$ which gives

$$\frac{\partial M}{\partial t_k} = \frac{\partial(S_1^{-1} h)}{\partial t_k} (h^{-1} S_2) + (S_1^{-1} h) \frac{\partial(h^{-1} S_2)}{\partial t_k}.$$

Using the commutation relation (3.6) and (3.10), we also have

$$\frac{\partial M}{\partial t_k} = M \Lambda^k = (S_1^{-1} h) (h^{-1} S_2) \Lambda^k.$$

As $A_2 = (S_2^T h^{-1})^{-1} \Lambda (S_2^T h^{-1})$, we obtain after some algebra

$$A_2^{-k} = \frac{\partial(S_1^{-1} h)^T}{\partial t_k} ((S_1^{-1} h)^T)^{-1} + (S_2^T h^{-1})^{-1} \frac{\partial(S_2^T h^{-1})}{\partial t_k}.$$

Since $(S_1^{-1} h)^T$ is upper triangular, the first term in the right hand side of this equation is upper triangular. As $S_2^T h^{-1}$ is lower triangular with all diagonal entries equal to 1, the second term is strictly lower triangular. Consequently, taking the strictly lower triangular part of both sides of the equation yields

$$(S_2^T h^{-1})^{-1} \frac{\partial(S_2^T h^{-1})}{\partial t_k} = (A_2^{-k})_{--},$$

which establishes (3.12), completing the proof. $\square$

We are now able to obtain a Lax pair representation for the Ablowitz-Ladik hierarchy.

Theorem 3.4. *The "dressed up" form of the moment equation (3.10) gives the following Lax pair representation for the Ablowitz-Ladik hierarchy on the semi-infinite CMV matrices (A_1, A_2)*

$$\boxed{\frac{\partial A_1}{\partial t_k} = [A_1, (A_1^k)_{--}], \quad \frac{\partial A_2}{\partial t_k} = [A_2, (A_2^{-k})_{--}], \quad \forall k \in \mathbb{Z}. \quad (3.13)}$$

Proof. As $A_1 = S_1 \Lambda S_1^{-1}$ and $A_2 = (S_2^T h^{-1})^{-1} \Lambda (S_2^T h^{-1})$, we have

$$\frac{\partial A_1}{\partial t_k} = \left[\frac{\partial S_1}{\partial t_k} S_1^{-1}, A_1 \right] \quad \text{and} \quad \frac{\partial A_2}{\partial t_k} = \left[A_2, (S_2^T h^{-1})^{-1} \frac{\partial(S_2^T h^{-1})}{\partial t_k} \right].$$

By Lemma 3.3 we obtain

$$\frac{\partial A_1}{\partial t_k} = [-(A_1^k)_{--}, A_1] \quad \text{and} \quad \frac{\partial A_2}{\partial t_k} = [A_2, (A_2^{-k})_{--}],$$

which establishes (3.13), concluding the proof. $\square$

Remark 3.5. Looking back at the explicit expressions for the entries of the CMV matrices in Theorem 2.8, the reader will observe that the entries of A_2 are obtained from those of A_1 by exchanging the roles of the variables x_n and y_n . Also A_1 contains as entries $-x_{2n+1}$ and y_{2n} and thus A_2 contains as entries x_{2n} and $-y_{2n+1}$, $n \geq 0$ (remember that $x_0 = y_0 = 1$). Thus the pair of Lax equations in (3.13) completely determines the Ablowitz-Ladik hierarchy in terms of the variables x_n and y_n .

Using the explicit expressions in terms of the variables x_n and y_n for the entries of the CMV matrices obtained in Theorem 2.8 and Theorem 3.4, one easily computes the equations for the vector fields T_1 and T_{-1}

$$\begin{aligned}\frac{\partial x_n}{\partial t_1} &= (1 - x_n y_n) x_{n+1}, & \frac{\partial x_n}{\partial t_{-1}} &= -(1 - x_n y_n) x_{n-1}, \\ \frac{\partial y_n}{\partial t_1} &= -(1 - x_n y_n) y_{n-1}, & \frac{\partial y_n}{\partial t_{-1}} &= (1 - x_n y_n) y_{n+1}.\end{aligned}$$

After the rescaling $x_n \rightarrow e^{-2t} x_n$, $y_n \rightarrow e^{2t} y_n$, the vector field $T_1 - T_{-1}$ writes

$$\begin{aligned}\dot{x}_n &= x_{n+1} - 2x_n + x_{n-1} - x_n y_n (x_{n+1} + x_{n-1}), \\ \dot{y}_n &= -y_{n+1} + 2y_n - y_{n-1} + x_n y_n (y_{n+1} + y_{n-1}),\end{aligned}$$

which are usually known as the Ablowitz-Ladik equations, see [1, 2]. Upon making the change of variable $t \rightarrow it$, when $y_n = \mp \overline{x_n}$ the system reduces to the equation

$$-i\dot{x}_n = x_{n+1} - 2x_n + x_{n-1} \pm |x_n|^2 (x_{n+1} + x_{n-1}),$$

which is a discrete version of the focusing/defocusing nonlinear Schrödinger equation.

3.2. A Lax pair for the master symmetries. In this section we translate the action of the master symmetries vector fields V_k , $k \in \mathbb{Z}$, defined on the bi-moments by (1.17), on the CMV matrices (A_1, A_2) .

We first decompose the vector fields V_k as follows

$$V_k = kT_k + \mathcal{V}_k, \quad (3.14)$$

where T_k are the Ablowitz-Ladik vector fields (3.9). At the level of the bi-moments, the vector fields $\mathcal{V}_k$ are given by

$$\boxed{\mathcal{V}_k \mu_j \equiv \frac{d}{du_k} \mu_j = j \mu_{j+k}, \quad j, k \in \mathbb{Z}.} \quad (3.15)$$

These vector fields satisfy the following commutation relations

$$\begin{aligned}[\mathcal{V}_k, \mathcal{V}_l] &= (l - k) \mathcal{V}_{k+l}, \\ [\mathcal{V}_k, T_l] &= l T_{k+l}.\end{aligned}$$

It follows that

$$[[\mathcal{V}_k, T_l], T_l] = 0, \quad \forall k, l \in \mathbb{Z}.$$

Consequently, like the vector fields V_k , the vector fields $\mathcal{V}_k$, $k \in \mathbb{Z}$, form a Virasoro algebra of master symmetries for the Ablowitz-Ladik hierarchy.

The differentiation of $\chi(z)$ with respect to z is defined by

$$\frac{d}{dz} \chi(z) = \delta \chi(z), \quad (3.16)$$

where

$$\delta = \Delta \Lambda^T, \quad \text{with } \Delta = \text{diag}(0, 1, -1, 2, -2, \dots), \quad (3.17)$$

and Λ is as in (3.5).

Remembering the notation (1.5), (3.15) writes

$$\frac{d}{du_k} \mu_{m,n} = (m - n) \mu_{m+k,n}$$

which is equivalent to the following equation on the bi-moment matrix M

$$\boxed{\frac{dM}{du_k} = \Delta \Lambda^k M - \Lambda^k M \Delta = [\Delta, \Lambda^k M].} \quad (3.18)$$

Remember from (3.2) that

$$f(z) = S_1 \chi(z), \quad g(z) = h (S_2^T)^{-1} \chi(z), \quad (3.19)$$

and, according to (2.6) and (2.7), these vectors satisfy

$$A_1 f(z) = z f(z), \quad A_1^T (h^{-1} g^*(z)) = z (h^{-1} g^*(z)), \quad (3.20)$$

$$A_2 g(z) = z g(z), \quad A_2^T (h^{-1} f^*(z)) = z (h^{-1} f^*(z)). \quad (3.21)$$

We define the semi-infinite matrices D_1, D_1^* and D_2, D_2^* by the relations

$$\frac{d}{dz} f(z) = D_1 f(z), \quad \frac{d}{dz} (h^{-1} g^*(z)) = (D_1^*)^T (h^{-1} g^*(z)), \quad (3.22)$$

$$\frac{d}{dz} g(z) = D_2 g(z), \quad \frac{d}{dz} (h^{-1} f^*(z)) = (D_2^*)^T (h^{-1} f^*(z)). \quad (3.23)$$

These matrices can be "dressed up" as explained in the next lemma.

Lemma 3.6. *We have*

$$D_1 = S_1 \Delta \Lambda^T S_1^{-1}, \quad D_1^* = -S_2 \Lambda^T \Delta S_2^{-1}, \quad (3.24)$$

$$D_2 = (S_2^T h^{-1})^{-1} \Delta \Lambda^T (S_2^T h^{-1}), \quad D_2^* = -(S_1^T h^{-1})^{-1} \Lambda^T \Delta (S_1^T h^{-1}), \quad (3.25)$$

with Δ as in (3.17).

Proof. Using (3.19) and (3.22), we have

$$D_1 f(z) = \frac{d}{dz} f(z) = S_1 \frac{d}{dz} \chi(z).$$

By definition of δ in (3.16) and (3.17), we get

$$D_1 f(z) = S_1 \delta S_1^{-1} f(z) = S_1 \Delta \Lambda^T S_1^{-1} f(z).$$

This proves the first formula in (3.24).

Using (3.19) and remembering from (2.5) that $g^*(z) = g(z^{-1})$, we have

$$\frac{d}{dz} g^*(z) = h (S_2^T)^{-1} \frac{d}{dz} \chi(z^{-1}) = -h (S_2^T)^{-1} z^{-2} \left(\frac{d}{du} \chi(u) \right) \Big|_{u=z^{-1}},$$

which gives, using (3.16), (3.17), (3.19) and remembering the definition (3.4) of the shift matrix Λ ,

$$\begin{aligned} \frac{d}{dz} g^*(z) &= -h (S_2^T)^{-1} \delta z^{-2} \chi(z^{-1}) \\ &= -h (S_2^T)^{-1} \Delta \Lambda^T \Lambda^2 \chi(z^{-1}) \\ &= -h (S_2^T)^{-1} \Delta \Lambda (h (S_2^T)^{-1})^{-1} g^*(z). \end{aligned}$$

Consequently, using the definition (3.22) of D_1^*

$$(D_1^*)^T (h^{-1} g^*(z)) = \frac{d}{dz} (h^{-1} g^*(z)) = -(S_2^T)^{-1} \Delta \Lambda S_2^T (h^{-1} g^*(z)).$$

This proves the second formula in (3.24).

The proof of (3.25) is identical to the proof of (3.24) using (3.19) and the definitions of D_2 and D_2^* in (3.23). This establishes the lemma. $\square$

Lemma 3.7. *We have for $k \in \mathbb{Z}$*

$$\frac{dS_1}{du_k} S_1^{-1} = -(D_1 A_1^{k+1})_{--} - (A_1^{k+1} D_1^*)_{--}, \quad (3.26)$$

$$(S_2^T h^{-1})^{-1} \frac{d(S_2^T h^{-1})}{du_k} = -(D_2 A_2^{1-k})_{--} - (A_2^{1-k} D_2^*)_{--}. \quad (3.27)$$

Proof. By substituting the factorisation $M = S_1^{-1} S_2$ of the moment matrix into (3.18), we obtain

$$-S_1^{-1} \frac{dS_1}{du_k} S_1^{-1} S_2 + S_1^{-1} \frac{dS_2}{du_k} = \Delta \Lambda^k S_1^{-1} S_2 - \Lambda^k S_1^{-1} S_2 \Delta.$$

Multiplying this equation on the left by S_1 and on the right by S_2^{-1} , we get

$$-\frac{dS_1}{du_k} S_1^{-1} + \frac{dS_2}{du_k} S_2^{-1} = \underbrace{S_1 \Delta \Lambda^k S_1^{-1}}_{\text{Term 1}} - \underbrace{S_1 \Lambda^k S_1^{-1} S_2 \Delta S_2^{-1}}_{\text{Term 2}}. \quad (3.28)$$

Using the factorisation of A_1 given in (3.7) and the factorisation of D_1 in (3.24), Term 1 gives

$$\text{Term 1} = S_1 \Delta \Lambda^T \Lambda^{k+1} S_1^{-1} = (S_1 \Delta \Lambda^T S_1^{-1}) (S_1 \Lambda^{k+1} S_1^{-1}) = D_1 A_1^{k+1}.$$

Similarly, the second term gives

$$\text{Term 2} = A_1^k S_2 \Delta S_2^{-1} = A_1^{k+1} A_1^{-1} S_2 \Delta S_2^{-1}.$$

Using the factorisation of A_1^{-1} in (3.8) we get

$$\text{Term 2} = A_1^{k+1} (S_2 \Lambda^T S_2^{-1}) S_2 \Delta S_2^{-1} = A_1^{k+1} (S_2 \Lambda^T \Delta S_2^{-1}) = -A_1^{k+1} D_1^*,$$

where we have used the expression of D_1^* in Lemma 3.6. Substituting these results in (3.28), we obtain

$$-\frac{dS_1}{du_k} S_1^{-1} + \frac{dS_2}{du_k} S_2^{-1} = D_1 A_1^{k+1} + A_1^{k+1} D_1^*.$$

The first term in the left-hand side is strictly lower triangular, while the second term in the left-hand side is upper triangular. Consequently, taking the strictly lower triangular part in both sides, we obtain

$$\frac{dS_1}{du_k} S_1^{-1} = -(D_1 A_1^{k+1})_{--} - (A_1^{k+1} D_1^*)_{--},$$

which establishes (3.26).

To establish the other formula, we substitute the factorisation $M = (S_1^{-1} h) (h^{-1} S_2)$ into equation (3.18) rewritten as

$$\frac{dM}{du_k} = [\Delta, M \Lambda^k],$$

which follows from the commutation relation (3.6). This gives

$$\begin{aligned} \frac{d(S_1^{-1} h)}{du_k} (h^{-1} S_2) + (S_1^{-1} h) \frac{d(h^{-1} S_2)}{du_k} \\ = \Delta (S_1^{-1} h) (h^{-1} S_2) \Lambda^k - (S_1^{-1} h) (h^{-1} S_2) \Lambda^k \Delta. \end{aligned}$$

Multiplying this equation on the left by $(S_1^{-1} h)^{-1}$ and on the right by $(h^{-1} S_2)^{-1}$, we get

$$\begin{aligned} (S_1^{-1} h)^{-1} \frac{d(S_1^{-1} h)}{du_k} + \frac{d(h^{-1} S_2)}{du_k} (h^{-1} S_2)^{-1} \\ = \underbrace{(S_1^{-1} h)^{-1} \Delta (S_1^{-1} h) (h^{-1} S_2) \Lambda^k (h^{-1} S_2)^{-1}}_{\text{Term 1}} - \underbrace{(h^{-1} S_2) \Lambda^k \Delta (h^{-1} S_2)^{-1}}_{\text{Term 2}}. \end{aligned} \quad (3.29)$$

Using the factorisation of A_2 in (3.7) and the factorisation of D_2 in (3.25), Term 2 gives

$$\begin{aligned} \text{Term 2} &= (h^{-1} S_2) \Lambda^{k-1} \Lambda \Delta (h^{-1} S_2)^{-1} \\ &= (h^{-1} S_2) \Lambda^{k-1} (h^{-1} S_2)^{-1} (h^{-1} S_2) \Lambda \Delta (h^{-1} S_2)^{-1} = (A_2^T)^{1-k} D_2^T. \end{aligned}$$

Similarly, using the factorisation of A_2 in (3.7), gives

$$\begin{aligned} \text{Term 1} &= (S_1^{-1} h)^{-1} \Delta (S_1^{-1} h) (A_2^T)^{-k} \\ &= (S_1^{-1} h)^{-1} \Delta (S_1^{-1} h) (A_2^T)^{-1} (A_2^T)^{1-k}. \end{aligned}$$

Using the factorisation of A_2^{-1} in (3.8) and the factorisation of D_2^* in (3.25), we get

$$\text{Term 1} = (h^{-1} S_1) \Delta \Lambda (h^{-1} S_1)^{-1} (A_2^T)^{1-k} = -(D_2^*)^T (A_2^T)^{1-k}.$$

Substituting these results in the transpose of (3.29), we obtain

$$\frac{d(S_1^{-1} h)^T}{du_k} ((S_1^{-1} h)^T)^{-1} + (S_2^T h^{-1})^{-1} \frac{d(S_2^T h^{-1})}{du_k} = -D_2 A_2^{1-k} - A_2^{1-k} D_2^*.$$

Since $(S_1^{-1}h)^T$ is upper triangular and $S_2^T h^{-1}$ is lower triangular with diagonal elements equal to 1, by taking the strictly lower part of both sides of this equation, we obtain (3.27). This concludes the proof of the lemma. $\square$

We are now able to obtain a Lax pair representation for the master symmetries vector fields $\mathcal{V}_k, k \in \mathbb{Z}$.

Theorem 3.8. *The "dressed up" form of the moment equation (3.18) gives the following Lax pair representation for the master symmetries vector fields $\mathcal{V}_k$ on the semi-infinite CMV matrices (A_1, A_2)*

$$\boxed{\begin{aligned} \frac{d}{du_k} A_1 &= \left[A_1, (D_1 A_1^{k+1})_{--} + (A_1^{k+1} D_1^*)_{--} \right], & \forall k \in \mathbb{Z}, \\ \frac{d}{du_k} A_2 &= \left[(D_2 A_2^{1-k})_{--} + (A_2^{1-k} D_2^*)_{--}, A_2 \right], & \forall k \in \mathbb{Z}, \end{aligned}} \quad (3.30)$$

or equivalently

$$\boxed{\begin{aligned} \frac{d}{du_k} A_1 &= A_1^{k+1} + \left[(D_1 A_1^{k+1})_+ - (A_1^{k+1} D_1^*)_{--}, A_1 \right], & \forall k \in \mathbb{Z}, \\ \frac{d}{du_k} A_2 &= A_2^{1-k} + \left[A_2, (A_2^{1-k} D_2^*)_+ - (D_2 A_2^{1-k})_{--} \right], & \forall k \in \mathbb{Z}. \end{aligned}}$$

Proof. As $A_1 = S_1 \Lambda S_1^{-1}$ and $A_2 = (S_2^T h^{-1})^{-1} \Lambda (S_2^T h^{-1})$, we have

$$\frac{dA_1}{du_k} = \left[\frac{dS_1}{du_k} S_1^{-1}, A_1 \right] \quad \text{and} \quad \frac{dA_2}{du_k} = \left[A_2, (S_2^T h^{-1})^{-1} \frac{d(S_2^T h^{-1})}{du_k} \right].$$

Using (3.26) and (3.27) in Lemma 3.7, we obtain (3.30).

From (3.20), (3.22) and from (3.21), (3.23), we deduce that $[A_1, D_1] = 1$ and $[D_2^*, A_2] = 1$. From these commutation relations, one readily obtains that

$$\begin{aligned} [A_1, (D_1 A_1^{k+1})_+] + [A_1, (D_1 A_1^{k+1})_{--}] &= A_1^{k+1}, \\ [(A_2^{1-k} D_2^*)_+, A_2] + [(A_2^{1-k} D_2^*)_{--}, A_2] &= A_2^{1-k}, \end{aligned}$$

which gives the equivalent formulation for the representation of the master symmetries on the CMV matrices (A_1, A_2) . This concludes the proof. $\square$

We notice that as a consequence of the Lax pair representation (3.13) for the AL hierarchy in Theorem 3.4, the relation between the vector fields V_k and $\mathcal{V}_k$ in (3.14) and the Lax pair representation (3.30) of $\mathcal{V}_k$ in Theorem 3.8, we have established the Lax pair representation (1.27), (1.28) of the vector fields V_k as announced in Theorem 1.1 in the Introduction.

Using the explicit form of the CMV matrices (A_1, A_2) in Theorem 2.8 and Theorem 3.8, remembering Remark 3.5, one can compute the first few master symmetries vector fields $\mathcal{V}_{-2}, \mathcal{V}_{-1}, \mathcal{V}_0, \mathcal{V}_1$ in terms of the variables x_n, y_n .

$$\begin{aligned}
\mathcal{V}_{-2}(x_n) &= (n-4)x_{n-2}(1-x_{n-1}y_{n-1})(1-x_ny_n) \\
&\quad - x_{n-1}(1-x_ny_n)((n-4)x_{n-1}y_n + (n-1)x_ny_{n+1}) \\
&\quad - 2x_{n-1}(1-x_ny_n) \sum_{k=1}^n y_k x_{k-1} + x_n \sum_{k=1}^n y_k^2 x_{k-1}^2 \\
&\quad - 2x_n \sum_{k=2}^n y_k x_{k-2} + 2x_n \sum_{k=2}^n y_k y_{k-1} x_{k-1} x_{k-2}, \\
\mathcal{V}_{-2}(y_n) &= -ny_{n+2}(1-x_ny_n)(1-x_{n+1}y_{n+1}) \\
&\quad + y_{n+1}(1-x_ny_n)(nx_ny_{n+1} + (n-1)x_{n-1}y_n) \\
&\quad + 2y_{n+1}(1-x_ny_n) \sum_{k=1}^n y_k x_{k-1} - y_n \sum_{k=1}^n y_k^2 x_{k-1}^2 \\
&\quad + 2y_n \sum_{k=2}^n y_k x_{k-2} - 2y_n \sum_{k=2}^n y_k y_{k-1} x_{k-1} x_{k-2}, \\
\mathcal{V}_{-1}(x_n) &= (n-2)x_{n-1}(1-x_ny_n) - x_n \sum_{k=1}^n y_k x_{k-1}, \\
\mathcal{V}_{-1}(y_n) &= -ny_{n+1}(1-x_ny_n) + y_n \sum_{k=1}^n y_k x_{k-1}, \\
\mathcal{V}_0(x_n) &= nx_n, \\
\mathcal{V}_0(y_n) &= -ny_n,
\end{aligned}$$

$$\begin{aligned}
\mathcal{V}_1(x_n) &= nx_{n+1}(1-x_ny_n) - x_n \sum_{k=1}^n x_k y_{k-1}, \\
\mathcal{V}_1(y_n) &= -(n-2)y_{n-1}(1-x_ny_n) + y_n \sum_{k=1}^n x_k y_{k-1}.
\end{aligned}$$

4. THE ACTION OF THE MASTER SYMMETRIES ON THE TAU-FUNCTIONS

As we recalled in the Introduction in formula (1.22), the tau-functions of the AL hierarchy are given by

$$\tau_n(t, s) = \det(\mu_{k-l}(t, s))_{0 \leq k, l < n}. \quad (4.1)$$

It immediately follows from the generating function of the elementary Schur polynomials (1.23) that

$$\frac{\partial}{\partial t_k} S_n(t) = S_{n-k}(t), \quad (4.2)$$

which shows that the formal solution of the AL hierarchy (1.16) on the moments is

$$\mu_j(t, s) = \sum_{m,n=0}^{\infty} S_m(t) S_n(s) \mu_{j+m-n}(0, 0), \quad \forall j \in \mathbb{Z}. \quad (4.3)$$

The expansion (1.24) of the tau-functions in terms of the Plücker coordinates (1.25) and the Schur polynomials (1.26) easily follows. Indeed, by substituting (4.3) into (4.1), we have

$$\begin{aligned} \tau_n(t, s) = & \sum_{\substack{0 \leq i_0, i_1, \dots, i_{n-1} \\ 0 \leq j_0, j_1, \dots, j_{n-1}}} \det [\mu_{k-l+i_k-j_l}(0, 0)]_{0 \leq k, l < n} \\ & \times S_{i_0}(t) \dots S_{i_{n-1}}(t) S_{j_0}(s) \dots S_{j_{n-1}}(s). \end{aligned}$$

Relabeling the indices as follows $i_k \mapsto i_k - k, j_l \mapsto j_l - l$, we get

$$\begin{aligned} \tau_n(t, s) &= \sum_{\substack{0 \leq i_0, \dots, i_{n-1} \\ 0 \leq j_0, \dots, j_{n-1}}} \det [\mu_{i_k-j_l}(0, 0)]_{0 \leq k, l < n} S_{i_0}(t) S_{i_1-1}(t) \dots S_{i_{n-1}-(n-1)}(t) \\ &\quad \times S_{j_0}(s) S_{j_1-1}(s) \dots S_{j_{n-1}-(n-1)}(s) \\ &= \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} \sum_{\sigma_1, \sigma_2 \in S_n} (-1)^{\sigma_1} (-1)^{\sigma_2} \det [\mu_{i_k-j_l}(0, 0)]_{0 \leq k, l < n} \\ &\quad \times S_{i_{\sigma_1(0)}}(t) S_{i_{\sigma_1(1)}-1}(t) \dots S_{i_{\sigma_1(n-1)}-(n-1)}(t) \\ &\quad \times S_{j_{\sigma_2(0)}}(s) S_{j_{\sigma_2(1)}-1}(s) \dots S_{j_{\sigma_2(n-1)}-(n-1)}(s) \\ &= \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s), \quad (4.4) \end{aligned}$$

with $(-1)^\sigma$ the sign of the permutation σ .

The aim of this section is to establish the second part of Theorem 1.1 in the Introduction.

Theorem 4.1. *For all $k \in \mathbb{Z}$, we have*

$$\begin{aligned} L_k^{(n)} \tau_n(t, s) = & \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} V_k \left(p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \right) S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s), \quad (4.5) \end{aligned}$$

with $L_k^{(n)}, k \in \mathbb{Z}$, defined as in (1.29), (1.30), (1.31), and $V_k \left(p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \right)$ the Lie derivative of the Plücker coordinates (1.25) in the direction of the master symmetries V_k of the AL hierarchy, as defined in (1.17).

This theorem is the key to the quick derivation of the various "Virasoro-type" constraints satisfied by *special* tau-functions of the AL hierarchy. As an illustration we establish the following result.

Corollary 4.2. *The partition function of the unitary matrix model*

$$\tau_n(t, s) = \int_{U(n)} \exp \left\{ \sum_{j=1}^{\infty} (t_j \operatorname{Tr} U^j + s_j \operatorname{Tr} U^{-j}) \right\} dU,$$

where $U(n)$ is the group of unitary $n \times n$ matrices and dU is the standard Haar measure, normalized so that the total volume is 1, satisfies the Virasoro constraints

$$L_k^{(n)} \tau_n(t, s) = 0, \quad \forall k \in \mathbb{Z},$$

with $L_k^{(n)}$ defined as in (1.29), (1.30) and (1.31).

Proof. This case corresponds to $\rho(z) = 1$ in the deformed weight (1.15), so that the initial moments (at time $(t, s) = (0, 0)$) are given by

$$\mu_j(0, 0) = \oint_{S^1} z^j \frac{dz}{2\pi i z} = \delta_{j,0},$$

with $\delta_{j,k}$ the usual Kronecker symbol. By the definition (1.17) of the master symmetries V_k , it follows that

$$V_k(\mu_j)|_{(t,s)=(0,0)} = (j+k)\mu_{j+k}(0,0) = (j+k)\delta_{j+k,0} = 0,$$

which, using the definition of the Plücker coordinates (1.25) and formula (4.5), establishes the result. $\square$

Remark 4.3. The unitary matrix model was first discussed in [24]. Corollary 4.2 was established by Haine and Vanderstichelen [21] in the more general case when the integration takes place over the unitary matrices whose spectrum miss an arc of the unit circle. Then the Virasoro constraints decouple into a time-part and a boundary-part. The method used in [21] was based on the standard trick of making a change of variable in the integral and exploiting the invariance of the integral. After [21] was completed, we noticed that Corollary 4.2 had already been obtained by Bowick, Morozov and Shevitz [8], using the same method. The authors obtain the equivalent constraints $(L_k^{(n)} - L_{-k}^{(n)})\tau_n(t, s) = 0, k \geq 1$, and $(L_k^{(n)} + L_{-k}^{(n)})\tau_n(t, s) = 0, k \geq 0$, but don't notice the commutation relations (1.32) of the centerless Virasoro algebra. Also in [4] Adler and van Moerbeke found the sl_2 subalgebra of constraints corresponding to $k = -1, 0, 1$, using the 2-Toda vertex operator.

Actually, in the proof of Theorem 4.1, we shall need to know that the operators $L_k^{(n)}, k \in \mathbb{Z}$, satisfy the commutation relations of the centerless Virasoro algebra. For the convenience of the reader we repeat the proof given in [21]. Consider the complex Lie algebra $\mathcal{A}$ given by the direct sum

of two commuting copies of the Heisenberg algebra with bases $\{\hbar_a, a_j | j \in \mathbb{Z}\}$ and $\{\hbar_b, b_j | j \in \mathbb{Z}\}$ and defining commutation relations

$$\begin{aligned} [\hbar_a, a_j] &= 0 \quad , \quad [a_j, a_k] = j\delta_{j,-k}\hbar_a, \\ [\hbar_b, b_j] &= 0 \quad , \quad [b_j, b_k] = j\delta_{j,-k}\hbar_b, \\ [\hbar_a, \hbar_b] &= 0 \quad , \quad [a_j, b_k] = 0 \quad , \quad [\hbar_a, b_j] = 0 \quad , \quad [\hbar_b, a_j] = 0, \end{aligned} \quad (4.6)$$

with $j, k \in \mathbb{Z}$. Let $\mathcal{B}$ be the space of formal power series in the variables $t_1, t_2, \dots$ and $s_1, s_2, \dots$, and consider the following representation of $\mathcal{A}$ in $\mathcal{B}$

$$\begin{aligned} a_j &= \frac{\partial}{\partial t_j} \quad , \quad a_{-j} = jt_j \quad , \quad b_j = \frac{\partial}{\partial s_j} \quad , \quad b_{-j} = js_j, \\ a_0 &= b_0 = \mu \quad , \quad \hbar_a = \hbar_b = 1, \end{aligned} \quad (4.7)$$

for $j > 0$, and $\mu \in \mathbb{C}$. Define the operators

$$A_k^{(n)} = \frac{1}{2} \sum_{j \in \mathbb{Z}} : a_{-j} a_{j+k} : \quad , \quad B_k^{(n)} = \frac{1}{2} \sum_{j \in \mathbb{Z}} : b_{-j} b_{j+k} :, \quad (4.8)$$

where $k \in \mathbb{Z}$, a_j, b_j are as in (4.7) with $\mu = n$, and where the colons indicate normal ordering, defined by

$$: a_j a_k : := \begin{cases} a_j a_k & \text{if } j \leq k, \\ a_k a_j & \text{if } j > k, \end{cases}$$

and a similar definition for $: b_j b_k :$, obtained by changing the a 's in b 's in the former. Expanding the expressions in (4.8) we obtain for $k > 0$

$$\begin{aligned} A_0^{(n)} &= \sum_{j>0} jt_j \frac{\partial}{\partial t_j} + \frac{n^2}{2}, \\ A_k^{(n)} &= \frac{1}{2} \sum_{0<j<k} \frac{\partial^2}{\partial t_j \partial t_{k-j}} + \sum_{j>k} (j-k)t_{j-k} \frac{\partial}{\partial t_j} + n \frac{\partial}{\partial t_k}, \\ A_{-k}^{(n)} &= \frac{1}{2} \sum_{0<j<k} j(k-j)t_j t_{k-j} + \sum_{j>k} jt_j \frac{\partial}{\partial t_{j-k}} + nkt_k, \end{aligned}$$

and similar expressions for $B_k^{(n)}$, by changing the t -variables in s -variables. Using these notations, we can rewrite (1.29), (1.30) and (1.31) as follows

$$\begin{aligned} L_k^{(n)} &= A_k^{(n)} - B_{-k}^{(n)} + \frac{1}{2} \sum_{j=1}^{k-1} (a_j - b_{-j})(a_{k-j} - b_{j-k}), \quad k \geq 1 \\ L_0^{(n)} &= A_0^{(n)} - B_0^{(n)}, \\ L_{-k}^{(n)} &= A_{-k}^{(n)} - B_k^{(n)} - \frac{1}{2} \sum_{j=1}^{k-1} (a_{-j} - b_j)(a_{j-k} - b_{k-j}), \quad k \geq 1. \end{aligned} \quad (4.9)$$

As shown in [22] (see Lecture 2) the operators $A_k^{(n)}$, $k \in \mathbb{Z}$, provide a representation of the Virasoro algebra in $\mathcal{B}$ with central charge $c = 1$, that

is

$$[A_k^{(n)}, A_l^{(n)}] = (k-l)A_{k+l}^{(n)} + \delta_{k,-l} \frac{k^3 - k}{12}, \quad (4.10)$$

for $k, l \in \mathbb{Z}$. Similarly, the operators $B_k^{(n)}$ satisfy the commutation relations

$$[B_k^{(n)}, B_l^{(n)}] = (k-l)B_{k+l}^{(n)} + \delta_{k,-l} \frac{k^3 - k}{12}, \quad (4.11)$$

for $k, l \in \mathbb{Z}$. Furthermore we have for $k, l \in \mathbb{Z}$

$$[a_k, A_l^{(n)}] = ka_{k+l}, \quad [b_k, B_l^{(n)}] = kb_{k+l}, \quad [a_k, B_l^{(n)}] = 0, \quad [b_k, A_l^{(n)}] = 0. \quad (4.12)$$

Proposition 4.4. *The operators $L_k^{(n)}$ defined as in (1.29), (1.30), (1.31) satisfy the commutation relations of the centerless Virasoro algebra*

$$[L_k^{(n)}, L_l^{(n)}] = (k-l)L_{k+l}^{(n)}, \quad \forall k, l \in \mathbb{Z}. \quad (4.13)$$

Proof. We give the proof for $k, l \geq 0$, the other cases being similar. As $[A_i^{(n)}, B_j^{(n)}] = 0$, $i, j \in \mathbb{Z}$, we have using (4.6), (4.10), (4.11) and (4.12)

$$\begin{aligned} [L_k^{(n)}, L_l^{(n)}] &= (k-l)(A_{k+l}^{(n)} - B_{-k-l}^{(n)}) - \frac{1}{2} \sum_{j=1}^{l-1} j(a_{j+k} - b_{-j-k})(a_{l-j} - b_{j-l}) \\ &\quad - \frac{1}{2} \sum_{j=1}^{l-1} (l-j)(a_j - b_{-j})(a_{k+l-j} - b_{j-k-l}) + \frac{1}{2} \sum_{j=1}^{k-1} j(a_{j+l} - b_{-j-l})(a_{k-j} - b_{j-k}) \\ &\quad + \frac{1}{2} \sum_{j=1}^{k-1} (k-j)(a_j - b_{-j})(a_{k+l-j} - b_{j-k-l}). \end{aligned}$$

Relabeling the indices in the sums, we have

$$\begin{aligned} [L_k^{(n)}, L_l^{(n)}] &= (k-l)(A_{k+l}^{(n)} - B_{-k-l}^{(n)}) \\ &\quad - \frac{1}{2} \sum_{j=k+1}^{k+l-1} (j-k)(a_j - b_{-j})(a_{k+l-j} - b_{j-k-l}) \\ &\quad - \frac{1}{2} \sum_{j=1}^{l-1} (l-j)(a_j - b_{-j})(a_{k+l-j} - b_{j-k-l}) \\ &\quad + \frac{1}{2} \sum_{j=l+1}^{k+l-1} (j-l)(a_j - b_{-j})(a_{k+l-j} - b_{j-k-l}) \\ &\quad + \frac{1}{2} \sum_{j=1}^{k-1} (k-j)(a_j - b_{-j})(a_{k+l-j} - b_{j-k-l}) \\ &= (k-l)L_{k+l}^{(n)}. \end{aligned}$$

This concludes the proof. $\square$

The plan of the rest of the section is as follows. After some algebraic preliminaries, we shall translate the master symmetries on the Plücker coordinates $p_{i_0, \dots, i_{n-1}}$. Next we shall compute the action of the Virasoro operators $S_{j_0, \dots, j_{n-1}}$ on the products of Schur polynomials $S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s)$. Finally we shall end with the proof of Theorem 4.1.

4.1. Some algebraic lemmas. We shall need the following lemmas. In order to formulate them, we introduce some notations. Given n vectors $x_1, \dots, x_n \in \mathbb{R}^n$, we shall denote by $|x_1 x_2 \dots x_n|$ the determinant of the $n \times n$ matrix formed with the columns x_i . Also, given two vectors x and y , $x \wedge y$ denotes the usual wedge product, with components $(x \wedge y)_{rs} = x_r y_s - x_s y_r$. Finally, for an $n \times n$ matrix A , A_r will denote the r th column of A , and A_r^T the r th column of the transposed matrix, and $\text{tr}(A)$ will mean the trace of A . With these conventions, we have the following lemma.

Lemma 4.5 (Haine-Semengue [20]). *Let A and B be $n \times n$ matrices, with A invertible. Then*

$$\begin{aligned} (i) \quad & \sum_{r=1}^n |A_1 \dots A_{r-1} B_r A_{r+1} \dots A_n| = (\det A) \text{tr}(B A^{-1}), \\ (ii) \quad & \sum_{1 \leq r < s \leq n} |A_1 \dots A_{r-1} B_r A_{r+1} \dots A_{s-1} B_s A_{s+1} \dots A_n| \\ & = (\det A) \sum_{1 \leq r < s \leq n} ((B A^{-1})_r \wedge (B A^{-1})_s)_{rs}. \end{aligned}$$

Proof. (i) Let A, B be $n \times n$ matrices, with A invertible. As A is invertible, its columns form a basis of $\mathbb{C}^n$ and thus we have

$$B_r = A c^{(r)} = \sum_j c_j^{(r)} A_j, \quad (4.14)$$

for a certain $c^{(r)} \in \mathbb{C}^n$, whose components are $c_j^{(r)} = (A^{-1} B)_{jr}$. It then follows that

$$\begin{aligned} \sum_{r=1}^n |A_1 \dots A_{r-1} B_r A_{r+1} \dots A_n| &= \sum_{r=1}^n |A_1 \dots A_{r-1} \left(\sum_j c_j^{(r)} A_j \right) A_{r+1} \dots A_n| \\ &= \det A \sum_{r=1}^n c_r^{(r)} \\ &= (\det A) \text{tr}(B A^{-1}). \end{aligned}$$

(ii) Using (4.14), we have

$$\begin{aligned}
& \sum_{1 \leq r < s \leq n} |A_1 \dots A_{r-1} B_r A_{r+1} \dots A_{s-1} B_s A_{s+1} \dots A_n| \\
&= \sum_{1 \leq r < s \leq n} |A_1 \dots A_{r-1} \left(\sum_j c_j^{(r)} A_j \right) A_{r+1} \dots \\
&\quad \quad \quad A_{s-1} \left(\sum_j c_j^{(s)} A_j \right) A_{s+1} \dots A_n| \\
&= \sum_{1 \leq r < s \leq n} |A_1 \dots A_{r-1} (c_r^{(r)} A_r + c_s^{(r)} A_s) A_{r+1} \dots \\
&\quad \quad \quad A_{s-1} (c_r^{(s)} A_r + c_s^{(s)} A_s) A_{s+1} \dots A_n| \\
&= \det A \sum_{1 \leq r < s \leq n} (c_r^{(r)} c_s^{(s)} - c_s^{(r)} c_r^{(s)}) \\
&= \det A \sum_{1 \leq r < s \leq n} ((A^{-1}B)_r \wedge (A^{-1}B)_s)_{rs}.
\end{aligned}$$

We thus obtain

$$\begin{aligned}
& \sum_{1 \leq r < s \leq n} |A_1 \dots A_{r-1} B_r A_{r+1} \dots A_{s-1} B_s A_{s+1} \dots A_n| \\
&= \det A \sum_{1 \leq r < s \leq n} ((BA^{-1})_r \wedge (BA^{-1})_s)_{rs},
\end{aligned}$$

where we have used the fact that for X, Y two $n \times n$ matrices, we have

$$\sum_{1 \leq r < s \leq n} ((XY)_r \wedge (XY)_s)_{rs} = \sum_{1 \leq r < s \leq n} ((YX)_r \wedge (YX)_s)_{rs}. \quad (4.15)$$

This concludes the proof of the lemma. $\square$

We will also need a transposed version of this lemma.

Lemma 4.6. *With the same conditions as in Lemma 4.5, we have*

$$\begin{aligned}
(i) \quad & \sum_{r=1}^n |A_1^T \dots A_{r-1}^T (B)_r^T A_{r+1}^T \dots A_n^T| = \sum_{r=1}^n |A_1 \dots A_{r-1} B_r A_{r+1} \dots A_n|, \\
(ii) \quad & \sum_{1 \leq r < s \leq n} |A_1^T \dots A_{r-1}^T B_r^T A_{r+1}^T \dots A_{s-1}^T B_s^T A_{s+1}^T \dots A_n^T| \\
&= \sum_{1 \leq r < s \leq n} |A_1 \dots A_{r-1} B_r A_{r+1} \dots A_{s-1} B_s A_{s+1} \dots A_n|.
\end{aligned}$$

Proof. Both formulas are direct consequences of Lemma 4.5, by observing that for X, Y two $n \times n$ matrices, we have (4.15) and

$$(X_r^T \wedge X_s^T)_{rs} = (X_r \wedge X_s)_{rs}.$$

$\square$

We give two consequences of this lemma. First we particularize the preceding lemma to the Plücker coordinates, and then we particularize it to the Schur polynomials.

Lemma 4.7. *For $m \in \mathbb{Z}$ we have*

$$\begin{aligned}
 (i) \quad \sum_{l=1}^n p_{j_0, \dots, j_{n-l-m}, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} &= \sum_{l=1}^n p_{j_0, \dots, j_{n-l}, \dots, j_{n-1}}^{i_0, \dots, i_{n-l}+m, \dots, i_{n-1}}, \\
 (ii) \quad \sum_{1 \leq r < s \leq n} p_{j_0, \dots, j_{n-s-m}, \dots, j_{n-r-m}, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \\
 &= \sum_{1 \leq r < s \leq n} p_{j_0, \dots, j_{n-s}, \dots, j_{n-r}, \dots, j_{n-1}}^{i_0, \dots, i_{n-s}+m, \dots, i_{n-r}+m, \dots, i_{n-1}}.
 \end{aligned}$$

Proof. Define the $n \times n$ matrices

$$A = (\mu_{i_k - j_l})_{0 \leq k, l \leq n-1}, \quad \text{and} \quad B(m) = (\mu_{i_k - j_l + m})_{0 \leq k, l \leq n-1}.$$

We then have

$$\begin{aligned}
 \sum_{l=1}^n p_{j_0, \dots, j_{n-l-m}, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} &= \sum_{l=1}^n |A_1 \dots A_{n-l-1} (B(m))_{n-l} A_{n-l+1} \dots A_{n-1}| \\
 &= \sum_{l=1}^n |A_1^T \dots A_{n-l-1}^T (B(m))_{n-l}^T A_{n-l+1}^T \dots A_{n-1}^T| \\
 &= \sum_{l=1}^n p_{j_0, \dots, j_{n-l}+m, \dots, j_{n-1}}^{i_0, \dots, i_{n-l}+m, \dots, i_{n-1}},
 \end{aligned}$$

where we have used Lemma 4.6(i) in the second equality. This proves (i). The proof of (ii) is similar. $\square$

Lemma 4.8. *The following holds*

$$\begin{aligned}
 (i) \quad \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) \\
 &= \det \begin{pmatrix} S_{i_{n-1}-n}(t) & S_{i_{n-1}-(n-2)}(t) & \cdots & S_{i_{n-1}}(t) \\ S_{i_{n-2}-n}(t) & S_{i_{n-2}-(n-2)}(t) & \cdots & S_{i_{n-2}}(t) \\ \vdots & \vdots & \ddots & \vdots \\ S_{i_0-n}(t) & S_{i_0-(n-2)}(t) & \cdots & S_{i_0}(t) \end{pmatrix}, \\
 (ii) \quad \sum_{l=1}^{n-1} S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)+1, \dots, i_1-1}(t) \\
 &= \det \begin{pmatrix} S_{i_{n-1}-(n-1)}(t) & \cdots & S_{i_{n-1}-2}(t) & S_{i_{n-1}}(t) \\ S_{i_{n-2}-(n-1)}(t) & \cdots & S_{i_{n-2}-2}(t) & S_{i_{n-2}}(t) \\ \vdots & \ddots & \vdots & \vdots \\ S_{i_1-(n-1)}(t) & \cdots & S_{i_1-2}(t) & S_{i_1}(t) \end{pmatrix}.
 \end{aligned}$$

Proof. We prove (i). Define the $n \times n$ matrices

$$A = (S_{i_{n-k}-(n-k)+l-k}(t))_{1 \leq k, l \leq n},$$

$$B(m) = (S_{i_{n-k}-(n-k)+l-k+m}(t))_{1 \leq k, l \leq n}.$$

We have $S_{i_{n-1}-(n-1), \dots, i_0}(t) = \det A$. It then follows that

$$\begin{aligned} & \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) \\ &= \sum_{l=1}^n |A_1^T \dots A_{n-l-1}^T (B(-1))_{n-l}^T A_{n-l+1}^T \dots A_{n-1}^T|. \end{aligned}$$

Using Lemma 4.6(i) we get

$$\begin{aligned} & \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) \\ &= \sum_{l=1}^n |A_1 \dots A_{n-l-1} (B(-1))_{n-l} A_{n-l+1} \dots A_{n-1}|. \end{aligned}$$

In the right-hand side, in the l^{th} term, the l^{th} and $(l-1)^{th}$ columns coincide in the determinant, provided that $l \neq 1$. Consequently, only the first term of the right-hand side gives a non zero contribution. This proves (i). The proof of (ii) is similar. $\square$

4.2. Expression of the master symmetries on the Plücker coordinates. We now translate the master symmetries on Plücker coordinates.

Lemma 4.9. *Let $V_k p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}}$ denote the Lie derivative of the Plücker coordinates in the direction of the vector fields V_k . Then for $k \in \mathbb{Z}$,*

$$\begin{aligned} & V_k p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \\ &= \sum_{l=0}^{n-1} (i_l + k) p_{i_0, \dots, i_{l-1}, i_l+k, i_{l+1}, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} - \sum_{l=0}^{n-1} j_l p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{l-1}, j_l-k, j_{l+1}, \dots, j_{n-1}} \\ &= \sum_{l=0}^{n-1} i_l p_{i_0, \dots, i_{l-1}, i_l+k, i_{l+1}, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} - \sum_{l=0}^{n-1} (j_l - k) p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{l-1}, j_l-k, j_{l+1}, \dots, j_{n-1}}. \end{aligned} \tag{4.16}$$

Proof. Fix $0 \leq i_0 < i_1 < \dots < i_{n-1}$ and $0 \leq j_0 < j_1 < \dots < j_{n-1}$. We introduce the $n \times n$ matrices

$$A = (\mu_{i_k-j_l}(0,0))_{0 \leq k, l \leq n-1}, \quad B(m) = (\mu_{i_k-j_l+m}(0,0))_{0 \leq k, l \leq n-1},$$

as well as the diagonal matrix $D = \text{diag}(j_0, \dots, j_{n-1})$. We notice that $p_{i_0, \dots, i_{n-1}} = \det A$, by definition of the Plücker coordinates. From the definition of V_k and using Leibniz's rule we find for $k \in \mathbb{Z}$

$$V_k p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} = \sum_{l=0}^{n-1} \begin{vmatrix} \mu_{i_0-j_0} & \mu_{i_0-j_1} & \cdots & \mu_{i_0-j_{n-1}} \\ \vdots & \vdots & & \vdots \\ \mu_{i_{l-1}-j_0} & \mu_{i_{l-1}-j_1} & \cdots & \mu_{i_{l-1}-j_{n-1}} \\ (i_l - j_0 + k)\mu_{i_l-j_0+k} & (i_l - j_1 + k)\mu_{i_l-j_1+k} & \cdots & (i_l - j_{n-1} + k)\mu_{i_l-j_{n-1}+k} \\ \mu_{i_{l+1}-j_0} & \mu_{i_{l+1}-j_1} & \cdots & \mu_{i_{l+1}-j_{n-1}} \\ \vdots & \vdots & & \vdots \\ \mu_{i_{n-1}-j_0} & \mu_{i_{n-1}-j_1} & \cdots & \mu_{i_{n-1}-j_{n-1}} \end{vmatrix},$$

or equivalently,

$$V_k p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} = \sum_{l=0}^{n-1} (i_l + k) p_{i_0, \dots, i_{l-1}, i_l+k, i_{l+1}, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} - \sum_{l=1}^n |A_1^T \cdots A_{l-1}^T (B(k)D)_l^T A_{l+1}^T \cdots A_n^T|.$$

Using Lemma 4.6(i) we obtain

$$\begin{aligned} V_k p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} &= \sum_{l=0}^{n-1} (i_l + k) p_{i_0, \dots, i_{l-1}, i_l+k, i_{l+1}, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \\ &\quad - \sum_{l=1}^n |A_1 \cdots A_{l-1} (B(k)D)_l A_{l+1} \cdots A_n| \\ &= \sum_{l=0}^{n-1} (i_l + k) p_{i_0, \dots, i_{l-1}, i_l+k, i_{l+1}, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \\ &\quad - \sum_{l=1}^n j_{l-1} |A_1 \cdots A_{l-1} (B(k))_l A_{l+1} \cdots A_n|. \end{aligned}$$

This gives the first equality in (4.16). The second equality in (4.16) can be derived from the first one by using Lemma 4.7(i). $\square$

4.3. Action of the Virasoro operators $L_k^{(n)}$ on the Schur polynomials. Next we shall compute the action of the Virasoro operators on the products of Schur polynomials $S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s)$. We have the following lemma.

Lemma 4.10.

$$\begin{aligned}
(i) \quad & L_0^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s) \\
&= \sum_{l=0}^{n-1} (i_l - j_l) S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s), \\
(ii) \quad & L_1^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s) \\
&= \sum_{l=1}^n i_{n-l} S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s) \\
&\quad - \sum_{l=1}^n (j_{n-l} + 1) S_{i_{n-1}-(n-1), \dots, i_0}(t) \\
&\quad \quad \times S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s), \\
(iii) \quad & L_2^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s) \\
&= \sum_{l=1}^n i_{n-l} S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-2, \dots, i_0}(t) \\
&\quad \quad \times S_{j_{n-1}-(n-1), \dots, j_0}(s) \\
&\quad + \sum_{1 \leq k < l \leq n} S_{i_{n-1}-(n-1), \dots, i_{n-k}-(n-k)-1, \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) \\
&\quad \quad \times S_{j_{n-1}-(n-1), \dots, j_0}(s) \\
&\quad - \sum_{l=1}^n (j_{n-l} + 2) S_{i_{n-1}-(n-1), \dots, i_0}(t) \\
&\quad \quad \times S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+2, \dots, j_0}(s) \\
&\quad - \sum_{1 \leq k < l \leq n} S_{i_{n-1}-(n-1), \dots, i_0}(t) \\
&\quad \quad \times S_{j_{n-1}-(n-1), \dots, j_{n-k}-(n-k)+1, \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s) \\
&\quad + s_1 \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s) \\
&\quad - s_1 \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s).
\end{aligned}$$

Proof. By using Leibniz's rule and (4.2) we have for $j \geq 1$,

$$\frac{\partial}{\partial t_j} S_{i_{n-1}-(n-1), \dots, i_0}(t) = \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-j, \dots, i_0}(t), \quad (4.17)$$

and

$$\begin{aligned} \frac{\partial^2}{\partial t_1^2} S_{i_{n-1}-(n-1), \dots, i_0}(t) &= \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-2, \dots, i_0}(t) \\ &+ 2 \sum_{1 \leq r < s \leq n} S_{i_{n-1}-(n-1), \dots, i_{n-r}-(n-r)-1, \dots, i_{n-s}-(n-s)-1, \dots, i_0}(t). \end{aligned} \quad (4.18)$$

Define the following $n \times n$ matrices

$$\begin{aligned} A(t) &:= \begin{pmatrix} S_{i_{n-1}-(n-1)}(t) & \dots & S_{i_{n-1}}(t) \\ \vdots & & \vdots \\ S_{i_0-(n-1)}(t) & \dots & S_{i_0}(t) \end{pmatrix}, \\ B(j, t) &:= \begin{pmatrix} S_{i_{n-1}-(n-1)-j}(t) & \dots & S_{i_{n-1}-j}(t) \\ \vdots & & \vdots \\ S_{i_0-(n-1)-j}(t) & \dots & S_{i_0-j}(t) \end{pmatrix}, \end{aligned}$$

and $D = \text{diag}(n-1, n-2, \dots, 0)$. We shall denote $\hat{A}(s)$ and $\hat{B}(j, s)$ the same matrices with $t \rightarrow s$ and $(i_0, \dots, i_{n-1}) \rightarrow (j_0, \dots, j_{n-1})$. From the definition (1.23) of the elementary Schur polynomials it follows easily that for $j \geq 0$,

$$\begin{aligned} \sum_{k=1}^{\infty} k t_k \frac{\partial}{\partial t_{k+j}} S_i(t) &= (i-j) S_{i-j}(t), \\ \sum_{k=j+1}^{\infty} k t_k \frac{\partial}{\partial t_{k-j}} S_i(t) &= (i+j) S_{i+j}(t) - \sum_{1 \leq l \leq j} l t_l S_{i+j-l}(t). \end{aligned}$$

Consequently, by first using Leibniz's rule and then Lemma 4.5(i) we have for $j \geq 0$

$$\begin{aligned} \sum_{k=1}^{\infty} k t_k \frac{\partial}{\partial t_{k+j}} S_{i_{n-1}-(n-1), \dots, i_0}(t) &= \sum_{l=1}^n (i_{n-l} - j) S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-j, \dots, i_0}(t) \\ &- (\det A(t)) \text{tr}(A(t)^{-1} B(j, t) D), \end{aligned} \quad (4.19)$$

$$\begin{aligned}
& \sum_{k=j+1}^{\infty} kt_k \frac{\partial}{\partial t_{k-j}} S_{i_{n-1}-(n-1), \dots, i_0}(t) \\
&= \sum_{l=1}^n (i_{n-l} + j) S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)+j, \dots, i_0}(t) \\
&\quad - (\det A(t)) \operatorname{tr}(A(t)^{-1} B(-j, t) D) \\
&\quad - \sum_{m=1}^j mt_m (\det A(t)) \operatorname{tr}(A(t)^{-1} B(m-j, t)). \tag{4.20}
\end{aligned}$$

We are now ready to prove the lemma.

(i) From (4.9), we have $L_0^{(n)} = A_0^{(n)} - B_0^{(n)}$. Using (4.19) with $j = 0$, we obtain

$$\begin{aligned}
& A_0^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) \\
&= \sum_{k=1}^{\infty} kt_k \frac{\partial}{\partial t_k} S_{i_{n-1}-(n-1), \dots, i_0}(t) + \frac{n^2}{2} S_{i_{n-1}-(n-1), \dots, i_0}(t) \\
&= \sum_{l=1}^n i_{n-l} S_{i_{n-1}-(n-1), \dots, i_0}(t) - (\det A(t)) \operatorname{tr}(A(t)^{-1} B(0, t) D) \\
&\quad + \frac{n^2}{2} S_{i_{n-1}-(n-1), \dots, i_0}(t).
\end{aligned}$$

We have $B(0, t) = A(t)$, and thus

$$\begin{aligned}
& (\det A(t)) \operatorname{tr}(A(t)^{-1} B(0, t) D) = (\det A(t)) \operatorname{tr}(D) \\
&= \frac{n(n-1)}{2} S_{i_{n-1}-(n-1), \dots, i_0}(t).
\end{aligned}$$

Consequently, we get

$$A_0^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) = \left[\sum_{l=1}^n i_{n-l} + \frac{n}{2} \right] S_{i_{n-1}-(n-1), \dots, i_0}(t).$$

Similarly, we get

$$B_0^{(n)} S_{j_{n-1}-(n-1), \dots, j_0}(s) = \left[\sum_{l=1}^n j_{n-l} + \frac{n}{2} \right] S_{j_{n-1}-(n-1), \dots, j_0}(s).$$

Combining both equations, we obtain (i).

(ii) From (4.9), we have $L_1^{(n)} = A_1^{(n)} - B_{-1}^{(n)}$. We compute, using (4.17) and (4.19)

$$\begin{aligned} A_1^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) &= \left[\sum_{j=1}^{\infty} j t_j \frac{\partial}{\partial t_{j+1}} + n \frac{\partial}{\partial t_1} \right] S_{i_{n-1}-(n-1), \dots, i_0}(t) \\ &= \sum_{l=1}^n (i_{n-l} + n - 1) S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) \\ &\quad - (\det A(t)) \operatorname{tr}(A(t)^{-1} B(1, t) D). \end{aligned}$$

By virtue of Lemma 4.5(i), we have

$$(\det A(t)) \operatorname{tr}(A(t)^{-1} B(1, t) D) = (n-1) |(B(1, t))_1 A_2(t) \dots A_n(t)|.$$

But by virtue of Lemma 4.8(i), this is equal to

$$\begin{aligned} (\det A(t)) \operatorname{tr}(A(t)^{-1} B(1, t) D) \\ = (n-1) \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t). \end{aligned}$$

Hence, we obtain

$$A_1^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) = \sum_{l=1}^n i_{n-l} S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t). \quad (4.21)$$

Similarly, we have using (4.20)

$$\begin{aligned} B_{-1}^{(n)} S_{j_{n-1}-(n-1), \dots, j_0}(s) &= \left[\sum_{j=2}^{\infty} j s_j \frac{\partial}{\partial s_{j-1}} + n s_1 \right] S_{j_{n-1}-(n-1), \dots, j_0}(s) \\ &= \sum_{l=1}^n (j_{n-l} + 1) S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s) \\ &\quad - (\det \hat{A}(s)) \operatorname{tr}(\hat{A}(s)^{-1} \hat{B}(-1, s) D) \\ &\quad - s_1 (\det \hat{A}(s)) \operatorname{tr}(\hat{A}(s)^{-1} \hat{B}(0, s)) + n s_1 S_{j_{n-1}-(n-1), \dots, j_0}(s). \end{aligned}$$

We have using Lemma 4.5(i)

$$(\det \hat{A}(s)) \operatorname{tr}(\hat{A}(s)^{-1} \hat{B}(-1, s) D) = 0,$$

and, obviously, we also have

$$(\det \hat{A}(s)) \operatorname{tr}(\hat{A}(s)^{-1} \hat{B}(0, s)) = n S_{j_{n-1}-(n-1), \dots, j_0}(s).$$

Consequently we obtain

$$\begin{aligned} B_{-1}^{(n)} S_{j_{n-1}-(n-1), \dots, j_0}(s) \\ = \sum_{l=1}^n (j_{n-l} + 1) S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s). \quad (4.22) \end{aligned}$$

Substracting (4.21) and (4.22) gives (ii).

(iii) From (4.9), we have

$$L_2^{(n)} = A_2^{(n)} - B_{-2}^{(n)} + \frac{1}{2} \left(\frac{\partial}{\partial t_1} - s_1 \right)^2.$$

We study separately the contributions of the three terms in the operator $L_2^{(n)}$ on the product of Schur functions. We start with the contribution of $A_2^{(n)}$. We compute, using (4.17), (4.18) and (4.19)

$$\begin{aligned} & A_2^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) \\ &= \left[\frac{1}{2} \frac{\partial^2}{\partial t_1^2} + \sum_{j=1}^{\infty} j t_j \frac{\partial}{\partial t_{j+2}} + n \frac{\partial}{\partial t_2} \right] S_{i_{n-1}-(n-1), \dots, i_0}(t) \\ &= \sum_{l=1}^n \left(i_{n-l} + n - \frac{3}{2} \right) S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-2, \dots, i_0}(t) \\ &\quad + \sum_{1 \leq k < l \leq n} S_{i_{n-1}-(n-1), \dots, i_{n-k}-(n-k)-1, \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) \\ &\quad - (\det A(t)) \operatorname{tr}(A(t)^{-1} B(2, t) D). \end{aligned}$$

The last term in this equation gives by developping the trace

$$\begin{aligned} & (\det A(t)) \operatorname{tr}(A(t)^{-1} B(2, t) D) \\ &= (\det A(t)) \left[(n-1) (A(t)^{-1} B(2, t))_{11} + (n-2) (A(t)^{-1} B(2, t))_{22} \right]. \end{aligned}$$

We have

$$\begin{aligned} & (\det A(t)) \operatorname{tr}(A(t)^{-1} B(2, t)) \\ &= (\det A(t)) \left[(A(t)^{-1} B(2, t))_{11} + (A(t)^{-1} B(2, t))_{22} \right], \end{aligned}$$

and by a short computation

$$(A(t)^{-1} B(2, t))_{22} = - \sum_{1 \leq k < l \leq n} \left((A(t)^{-1} B(1, t))_k \wedge (A(t)^{-1} B(1, t))_l \right)_{kl}.$$

Consequently we have

$$\begin{aligned} & (\det A(t)) \operatorname{tr}(A(t)^{-1} B(2, t) D) = (n-1) (\det A(t)) \operatorname{tr}(A(t)^{-1} B(2, t)) \\ &\quad + (\det A(t)) \sum_{1 \leq k < l \leq n} \left((A(t)^{-1} B(1, t))_k \wedge (A(t)^{-1} B(1, t))_l \right)_{kl}. \end{aligned}$$

Using Lemma 4.5, we obtain

$$\begin{aligned}
& (\det A(t)) \operatorname{tr}(A(t)^{-1} B(2, t) D) \\
&= (n-1) \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-2, \dots, i_0}(t) \\
&+ \sum_{1 \leq k < l \leq n} S_{i_{n-1}-(n-1), \dots, i_{n-k}-(n-k)-1, \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t).
\end{aligned}$$

Hence, we get

$$\begin{aligned}
A_2^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) \\
= \sum_{l=1}^n \left(i_{n-l} - \frac{1}{2}\right) S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-2, \dots, i_0}(t). \quad (4.23)
\end{aligned}$$

We now turn to the contribution of $B_{-2}^{(n)}$. We have using (4.20)

$$\begin{aligned}
& B_{-2}^{(n)} S_{j_{n-1}-(n-1), \dots, j_0}(s) \\
&= \left[\frac{1}{2} s_1^2 + \sum_{j=3}^{\infty} j s_j \frac{\partial}{\partial s_{j-2}} + 2n s_2 \right] S_{j_{n-1}-(n-1), \dots, j_0}(s) \\
&= \left[\frac{1}{2} s_1^2 + 2n s_2 \right] S_{j_{n-1}-(n-1), \dots, j_0}(s) \\
&+ \sum_{l=1}^n (j_{n-l} + 2) S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+2, \dots, j_0}(s) \\
&- (\det \hat{A}(s)) \operatorname{tr}(\hat{A}(s)^{-1} \hat{B}(-2, s) D) \\
&- \sum_{m=1}^2 m s_m (\det \hat{A}(s)) \operatorname{tr}(\hat{A}(s)^{-1} \hat{B}(m-2, s)).
\end{aligned}$$

By a similar argument as above, we have

$$\begin{aligned}
& (\det \hat{A}(s)) \operatorname{tr}(\hat{A}(s)^{-1} \hat{B}(-2, s) D) \\
&= -(\det \hat{A}(s)) \sum_{1 \leq k < l \leq n} \left((\hat{A}(s)^{-1} \hat{B}(-1, s))_k \wedge (\hat{A}(s)^{-1} \hat{B}(-1, s))_l \right)_{kl},
\end{aligned}$$

and thus using Lemma 4.5(ii), we obtain

$$\begin{aligned}
& (\det \hat{A}(s)) \operatorname{tr}(\hat{A}(s)^{-1} \hat{B}(-2, s) D) \\
&= - \sum_{1 \leq k < l \leq n} S_{j_{n-1}-(n-1), \dots, j_{n-k}-(n-k)+1, \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s).
\end{aligned}$$

We also have, using Lemma 4.5(i),

$$\begin{aligned} & \sum_{m=1}^2 m s_m (\det \hat{A}(s)) \operatorname{tr} (\hat{A}(s)^{-1} \hat{B}(m-2, s)) \\ &= s_1 \sum_{l=1}^n S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s) + 2n s_2 S_{j_{n-1}-(n-1), \dots, j_0}(s). \end{aligned}$$

Consequently, we have

$$\begin{aligned} & B_{-2}^{(n)} S_{j_{n-1}-(n-1), \dots, j_0}(s) \\ &= \sum_{l=1}^n (j_{n-l} + 2) S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+2, \dots, j_0}(s) \\ & \quad + \sum_{1 \leq k < l \leq n} S_{j_{n-1}-(n-1), \dots, j_{n-k}-(n-k)+1, \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s) \\ & \quad - s_1 \sum_{l=1}^n S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s) \\ & \quad + \frac{1}{2} s_1^2 S_{j_{n-1}-(n-1), \dots, j_0}(s). \end{aligned} \tag{4.24}$$

Finally, we turn to the contribution of the term $\frac{1}{2} \left(\frac{\partial}{\partial t_1} - s_1 \right)^2$. We have using (4.17) and (4.18)

$$\begin{aligned} & \frac{1}{2} \left[\frac{\partial}{\partial t_1} - s_1 \right]^2 S_{i_{n-1}-(n-1), \dots, i_0}(t) \\ &= \frac{1}{2} \left[\frac{\partial^2}{\partial t_1^2} - 2s_1 \frac{\partial}{\partial t_1} + s_1^2 \right] S_{i_{n-1}-(n-1), \dots, i_0}(t) \\ &= \frac{1}{2} \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-2, \dots, i_0}(t) \\ & \quad + \sum_{1 \leq k < l \leq n} S_{i_{n-1}-(n-1), \dots, i_{n-k}-(n-k)-1, \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) \\ & \quad - s_1 \sum_{l=1}^n S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) + \frac{1}{2} s_1^2 S_{i_{n-1}-(n-1), \dots, i_0}(t). \end{aligned} \tag{4.25}$$

Combining (4.23), (4.24) and (4.25), we obtain (iii). $\square$

Remark 4.11. We observe that by definition of the operators $L_k^{(n)}$ we have

$$\begin{aligned} & L_{-k}^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s) \\ &= -L_k^{(n)} S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_0}(s) \Big|_{(i_0, \dots, i_{n-1}) \leftrightarrow (j_0, \dots, j_{n-1})}^{t \leftrightarrow s}. \end{aligned}$$

4.4. Proof of the main theorem. We now turn to the last part of this section. We will prove Theorem 4.1. We first prove the following lemma.

Lemma 4.12.

$$\begin{aligned}
& \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} S_{i_{n-1}-(n-1), \dots, i_0}(t) \\
& \quad \times S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s) \\
& + \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 < j_1 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{-1, j_1, \dots, j_{n-1}} S_{i_{n-1}-(n-1), \dots, i_0}(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 0}(s) \\
& = \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) \\
& \quad \times S_{j_{n-1}-(n-1), \dots, j_0}(s). \tag{4.26}
\end{aligned}$$

Proof. For simplicity, we will use the notations

$$\mathcal{S}_i(t) = S_{i_{n-1}-(n-1), \dots, i_0}(t), \quad \mathcal{S}_j(s) = S_{j_{n-1}-(n-1), \dots, j_0}(s), \tag{4.27}$$

when no 'special' shift on the indices of the Schur functions occur. Relabeling each term in the first sum of the left-hand side of (4.26) in the following way $j_{n-l} \mapsto j_{n-l} - 1$ gives

$$\begin{aligned}
& \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s) \\
& = \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-l}-1 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-l}-1, \dots, j_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s).
\end{aligned}$$

On the one hand, for a fixed $1 \leq l \leq n-1$, if $j_{n-l} = j_{n-l-1} + 1$, then $p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-l}-1, \dots, j_{n-1}} = 0$. On the other hand, for a fixed $2 \leq l \leq n$, if $j_{n-l} =$

j_{n-l+1} , then $S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l), \dots, j_0}(s) = 0$. Therefore

$$\begin{aligned} & \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s) \\ &= \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s) \\ &\quad - \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 < j_1 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 0}(s). \end{aligned}$$

Consequently, the left-hand side of (4.26) is equal to

$$\begin{aligned} & \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s) \\ &+ \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 < j_1 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 0}(s) \\ &= \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s). \end{aligned} \quad (4.28)$$

Similarly, one can show that the right-hand side of (4.26) is equal to

$$\begin{aligned} & \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}} \mathcal{S}_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) \mathcal{S}_j(s) \\ &= \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-l}+1, \dots, i_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s). \end{aligned} \quad (4.29)$$

By virtue of Lemma 4.7(i), (4.28) and (4.29) are equal. $\square$

Proof of Theorem 4.1: We will prove the theorem for $k \geq 0$. The case $k < 0$ is similar. Using the Plücker expansion (4.4) of $\tau_n(t)$, and Lemmas 4.9 and 4.10 we have for $k = 0, 1$, using the notations (4.27),

$$\begin{aligned} V_k \tau_n(s, t) &= \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} V_k p_{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s) \\ &= \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}} L_k^{(n)} \mathcal{S}_i(t) \mathcal{S}_j(s) \\ &= L_k^{(n)} \tau_n(s, t), \end{aligned}$$

where, in the second equality, we have performed some relabeling of the indices as in the proof of Lemma 4.12. We will finish the proof with the case $k = 2$, for which we provide some more details, but first we prove the theorem for general $k \geq 3$. We proceed by induction. Assume the theorem holds for some $k \geq 2$. We will establish it for $k + 1$. The argument follows from the commutation relations (4.13) and (1.18). We have

$$\begin{aligned}
(k-1)V_{k+1}\tau_n(s, t) &= \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} [V_1, V_k] p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s) \\
&= \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} [L_k^{(n)}, L_1^{(n)}] \mathcal{S}_i(t) \mathcal{S}_j(s) \\
&= (k-1)L_{k+1}^{(n)} \tau_n(s, t),
\end{aligned}$$

where in the second equality we have used the induction hypothesis.

We now provide some details for the case $k = 2$. Using Lemmas 4.10 and 4.12 we have

$$\begin{aligned}
L_2^{(n)} \tau_n(s, t) &= T_1 + T_2 + T_3 + T_4 \\
&\quad - s_1 \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 < j_1 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{-1, j_1, \dots, j_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 0}(s), \quad (4.30)
\end{aligned}$$

with

$$\begin{aligned}
T_1 &:= \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \sum_{l=1}^n i_{n-l} S_{i_{n-1}-(n-1), \dots, i_{n-l}-(n-l)-2, \dots, i_0}(t) \mathcal{S}_j(s), \\
T_2 &:= - \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \\
&\quad \times \sum_{l=1}^n (j_{n-l} + 2) \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+2, \dots, j_0}(s), \\
T_3 &:= \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \\
&\quad \times \sum_{1 \leq k < l \leq n} S_{i_{n-1}-(n-1), \dots, i_{n-k}-(n-k)-1, \dots, i_{n-l}-(n-l)-1, \dots, i_0}(t) \mathcal{S}_j(s), \\
T_4 &:= - \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \\
&\quad \times \sum_{1 \leq k < l \leq n} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_{n-k}-(n-k)+1, \dots, j_{n-l}-(n-l)+1, \dots, j_0}(s).
\end{aligned}$$

We will consider separately the four terms T_1, T_2, T_3, T_4 . By arguments similar to that used in the proof of Lemma 4.12, and using the fact that $S_{i_{n-1}-(n-1), \dots, i_0}(t) = 0$ if $i_k < 0$ for some $0 \leq k \leq n-1$, we get for T_1

$$\begin{aligned} T_1 &= \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} (i_{n-l} + 2) p_{i_0, \dots, i_{n-l}+2, \dots, i_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s) \\ &+ \sum_{l=1}^{n-1} \sum_{\substack{-1 \leq i_0-1 < \dots < i_{n-l-1}-1 \\ = i_{n-l} < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} (i_{n-l} + 2) p_{i_0, \dots, i_{n-l}+2, \dots, i_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s) \\ &- \sum_{l=2}^n \sum_{\substack{-1 \leq i_0-1 < \dots < i_{n-l-1}-1 \\ < i_{n-l}+1 = i_{n-l+1} < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} (i_{n-l} + 2) p_{i_0, \dots, i_{n-l}+2, \dots, i_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s). \end{aligned}$$

The two last terms in this expression annihilate, i.e.

$$\begin{aligned} 0 &= \sum_{l=1}^{n-1} \sum_{\substack{-1 \leq i_0-1 < \dots < i_{n-l-1}-1 \\ = i_{n-l} < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} (i_{n-l} + 2) p_{i_0, \dots, i_{n-l}+2, \dots, i_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s) \\ &- \sum_{l=2}^n \sum_{\substack{-1 \leq i_0-1 < \dots < i_{n-l-1}-1 \\ < i_{n-l}+1 = i_{n-l+1} < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} (i_{n-l} + 2) p_{i_0, \dots, i_{n-l}+2, \dots, i_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s). \end{aligned} \tag{4.31}$$

Indeed, we have for $1 \leq l \leq n-1$

$$\begin{aligned} &\sum_{\substack{-1 \leq i_0-1 < \dots < i_{n-l-1}-1 \\ = i_{n-l} < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} (i_{n-l} + 2) p_{i_0, \dots, i_{n-l}+2, \dots, i_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s) = \\ &\sum_{\substack{-1 \leq k_0-1 < \dots < k_{n-l-1}-1 \\ = k_{n-l}-1 < \dots < k_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} (k_{n-l-1} + 2) p_{k_0, \dots, k_{n-l-2}, k_{n-l}, k_{n-l-1}+2, k_{n-l+1}, \dots, k_{n-1}} \times \end{aligned}$$

$$S_{k_{n-1}-(n-1), \dots, k_{n-l+1}-(n-l+1), k_{n-l-1}-(n-l), k_{n-l}-(n-l-1), k_{n-l-2}-(n-l-2), \dots, k_0}(t) \mathcal{S}_j(s),$$

where we have made the relabeling $i_{n-l-1} \mapsto k_{n-l}$, $i_{n-l} \mapsto k_{n-l-1}$, and $i_m \mapsto k_m$ if $m \neq n-l-1, n-l$. As the Plücker coordinates and the Schur functions are determinants, we have, permuting lines in the determinants,

$$p_{k_0, \dots, k_{n-l-2}, k_{n-l}, k_{n-l-1}+2, k_{n-l+1}, \dots, k_{n-1}} = -p_{k_0, \dots, k_{n-l-1}+2, \dots, k_{n-1}},$$

and

$$\begin{aligned} S_{k_{n-1}-(n-1), \dots, k_{n-l+1}-(n-l+1), k_{n-l-1}-(n-l), k_{n-l}-(n-l-1), k_{n-l-2}-(n-l-2), \dots, k_0}(t) \\ = -S_k(t), \end{aligned}$$

and hence

$$\begin{aligned} \sum_{\substack{-1 \leq i_0-1 < \dots < i_{n-l-1}-1 = i_{n-l} < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} (i_{n-l} + 2) p_{i_0, \dots, i_{n-l}+2, \dots, i_{n-1}} S_i(t) S_j(s) = \\ \sum_{\substack{-1 \leq k_0-1 < \dots < k_{n-l-1} = k_{n-l}-1 < \dots < k_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} (k_{n-l-1} + 2) p_{k_0, \dots, k_{n-l-1}+2, \dots, k_{n-1}} S_k(t) S_j(s). \end{aligned}$$

Summing this expression for $1 \leq l \leq n-1$, and relabeling $l \mapsto l-1$ we get (4.31). Consequently we obtain

$$T_1 = \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} (i_{n-l} + 2) p_{i_0, \dots, i_{n-l}+2, \dots, i_{n-1}} S_i(t) S_j(s). \quad (4.32)$$

By similar arguments, we have

$$\begin{aligned} T_2 = - \sum_{l=1}^n \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} j_{n-l} p_{i_0, \dots, i_{n-1}} S_i(t) S_j(s) \\ + \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-1}} S_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 1}(s), \end{aligned} \quad (4.33)$$

$$T_3 = \sum_{1 \leq k < l \leq n} \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-l}+1, \dots, i_{n-k}+1, \dots, i_{n-1}} S_i(t) S_j(s), \quad (4.34)$$

$$\begin{aligned} T_4 = - \sum_{1 \leq k < l \leq n} \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} p_{i_0, \dots, i_{n-1}} S_i(t) S_j(s) \\ + \sum_{1 \leq k \leq n-1} \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 < j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-k}-1, \dots, j_{n-1}} S_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 0}(s). \end{aligned} \quad (4.35)$$

Substituting (4.32), (4.33), (4.34) and (4.35) in (4.30), using Lemma 4.7(ii) and Lemma 4.9 we obtain

$$\begin{aligned}
L_2^{(n)} \tau_n(s, t) = & \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} V_2 p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s) \\
& + \sum_{k=1}^{n-1} \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-k-1}, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 0}(s) \\
& - s_1 \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 0}(s) \\
& + \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 1}(s).
\end{aligned}$$

We prove that the last three terms in this expression annihilate

$$\begin{aligned}
0 = & \sum_{k=1}^{n-1} \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-k-1}, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 0}(s) \\
& - s_1 \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 0}(s) \\
& + \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 1}(s), \tag{4.36}
\end{aligned}$$

and hence

$$L_2^{(n)} \tau_n(s, t) = \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_0 < \dots < j_{n-1}}} V_2 p_{i_0, \dots, i_{n-1}}^{j_0, \dots, j_{n-1}} \mathcal{S}_i(t) \mathcal{S}_j(s). \tag{4.37}$$

Indeed, developping the determinant $S_{j_{n-1}-(n-1), \dots, j_1-1, 1}(s)$ with respect to the last line, using the fact that the first elementary Schur polynomials are $S_0(s) = 1$ and $S_1(s) = s_1$, and Lemma 4.8(ii), we have

$$\begin{aligned}
& \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 1}(s) \\
= & s_1 \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1}(s) \\
& - \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) \sum_{l=1}^{n-1} S_{j_{n-1}-(n-1), \dots, j_{n-l}-(n-l)+1, \dots, j_1-1}(s).
\end{aligned}$$

By an argument similar to that of the proof of Lemma 4.12, we get

$$\begin{aligned}
& \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1, 1}(s) \\
&= s_1 \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1}(s) \\
&\quad - \sum_{l=1}^{n-1} \sum_{\substack{0 \leq i_0 < \dots < i_{n-1} \\ 0 \leq j_1 < \dots < j_{n-1}}} p_{-1, j_1, \dots, j_{n-l}-1, \dots, j_{n-1}}^{i_0, \dots, i_{n-1}} \mathcal{S}_i(t) S_{j_{n-1}-(n-1), \dots, j_1-1}(s).
\end{aligned}$$

Noticing that $S_{j_{n-1}-(n-1), \dots, j_1-1}(s) = 0$ when $j_1 = 0$, and

$$S_{j_{n-1}-(n-1), \dots, j_1-1}(s) = S_{j_{n-1}-(n-1), \dots, j_1-1, 0}(s),$$

when $j_1 > 0$, we get (4.36), and hence (4.37). This proves the case $k = 2$ and finishes the proof. $\square$

REFERENCES

- [1] M. J. Ablowitz, J. F. Ladik, *Nonlinear differential-difference equations*, J. Math. Phys. **16** (1975) 598-603.
- [2] M. J. Ablowitz, J. F. Ladik, *Nonlinear differential-difference equations and Fourier analysis*, J. Math. Phys. **17** (1976) 1011-1018.
- [3] M. Adler, P. van Moerbeke, *Compatible Poisson structures and the Virasoro algebra*, Comm. Pure Appl. Math. **47** (1994) 5-37.
- [4] M. Adler, P. van Moerbeke, *Integrals over classical groups, random permutations, Toda and Toeplitz lattices*, Comm. Pure Appl. Math. **54** (2001) 153-205.
- [5] M. Adler, P. van Moerbeke, *Recursion relations for unitary integrals, combinatorics and the Toeplitz lattice*, Comm. Math. Phys. **237** (2003) 397-440.
- [6] D. Aldous, P. Diaconis, *Longest increasing subsequences: from patience sorting to the Baik-Deift-Johansson theorem*, Bull. Amer. Math. Soc. **36** (1999) 413-432.
- [7] A. Borodin, *Discrete gap probabilities and discrete Painlevé equations*, Duke Math. J. **117** (2003) 489-542.
- [8] M. J. Bowick, A. Morozov, D. Shevitz, *Reduced unitary matrix models and the hierarchy of τ -functions*, Nucl. Phys. B **354** (1991) 496-530.
- [9] M. J. Cantero, L. Moral, L. Velázquez, *Five-diagonal matrices and zeros of orthogonal polynomials on the unit circle*, Linear Algebra Appl. **362** (2003) 29-56.
- [10] R. Cruz-Barroso, P. González-Vera, *A Christoffel-Darboux formula and a Favard's theorem for orthogonal Laurent polynomials on the unit circle*, J. Comput. Appl. Math. **179** (2005) 157-173.
- [11] P. A. Damianou, *Master symmetries and R-matrices for the Toda lattice*, Lett. Math. Phys. **20** (1990) 101-112.
- [12] P. A. Damianou, *Multiple hamiltonian structures for Toda-type systems*, J. Math. Phys. **35** (1994) 5511-5541.
- [13] J. J. Duistermaat, F. A. Grünbaum, *Differential equations in the spectral parameter*, Comm. Math. Phys. **103** (1986) 177-240.
- [14] L. Faybusovich, M. Gekhtman, *Poisson brackets on rational functions and multi-Hamiltonian structure for integrable lattices*, Phys. Lett. A **272** (2000) 236-244.

- [15] P. J. Forrester, N. S. Witte, *Bi-orthogonal polynomials on the unit circle, regular semi-classical weights and integrable systems*, Constr. Approx. **24** (2006) 201-237.
- [16] B. Fuchssteiner, *Mastersymmetries, higher order time-dependent symmetries and conserved densities of nonlinear evolution equations*, Progr. Theor. Phys. **70** (1983) 1508-1522.
- [17] F. Gesztesy, H. Holden, J. Michor, G. Teschl, *Soliton Equations and Their Algebro-Geometric Solutions. Vol. 2. (1+1)-Dimensional Discrete Models*, Cambridge studies in advanced mathematics 114, Cambridge: Cambridge University Press, 2008.
- [18] F.A. Grünbaum, L. Haine, *A theorem of Bochner, revisited*, in Algebraic aspects of integrable systems, Progr. Nonlinear Differential Equations Appl. **26** 143-172, Birkhäuser Boston, Boston, MA, 1997.
- [19] L. Haine, E. Horozov, *Toda orbits of Laguerre polynomials and representations of the Virasoro algebra*, Bull. Sci. Math. **117** (1993) 485-518.
- [20] L. Haine, J. P. Semengue, *The Jacobi polynomial ensemble and the Painlevé VI equation*, J. Math. Phys. **40** (1999) 2117-2134.
- [21] L. Haine, D. Vanderstichelen, *A centerless representation of the Virasoro algebra associated with the unitary circular ensemble*, J. Comput. Appl. Math. **236** (2011) 19-27.
- [22] V. G. Kac, A. K. Raina, *Bombay Lectures on Highest Weight Representations of Infinite Dimensional Lie Algebras*, Adv. Series in Math. Phys., Vol.2, World Scientific, Singapore, 1987.
- [23] F. Magri, J. P. Zubelli, *Differential equations in the spectral parameter, Darboux transformations and a hierarchy of master symmetries for KdV*, Comm. Math. Phys. **141** (1991) 329-351.
- [24] E. J. Martinec, *On the origin of integrability in matrix models*, Comm. Math. Phys. **138** (1991) 437-449.
- [25] I. Nenciu, *Lax pair for the Ablowitz-Ladik system via orthogonal polynomials on the unit circle*, Int. Math. Res. Not. **11** (2005) 647-686.
- [26] E. M. Rains, *Increasing subsequences and the classical groups*, Electron. J. Combinatorics, **5**:R12 (1998).
- [27] B. Simon, *Orthogonal Polynomials on the Unit Circle, Part 1: Classical Theory*. Providence, RI: Amer. Math. Soc. (2005).
- [28] B. Simon, *Orthogonal Polynomials on the Unit Circle, Part 2: Spectral Theory*. Providence, RI: Amer. Math. Soc. (2005).
- [29] C. A. Tracy, H. Widom, *Fredholm determinants, differential equations and matrix models*, Comm. Math. Phys. **163** (1994) 33-72.
- [30] D. Vanderstichelen, Ph. D. thesis, Université catholique de Louvain, 2011.

LUC HAINE, INSTITUT DE RECHERCHE EN MATHÉMATIQUE ET PHYSIQUE, UNIVERSITÉ CATHOLIQUE DE LOUVAIN, CHEMIN DU CYCLOTRON 2, 1348 LOUVAIN-LA-NEUVE, BELGIUM, CORRESPONDING AUTHOR, TEL.:+32 10 47 3162; FAX:+32 10 47 2530.

E-mail address: luc.haine@uclouvain.be

DIDIER VANDERSTICHELEN, INSTITUT DE RECHERCHE EN MATHÉMATIQUE ET PHYSIQUE, UNIVERSITÉ CATHOLIQUE DE LOUVAIN, CHEMIN DU CYCLOTRON 2, 1348 LOUVAIN-LA-NEUVE, BELGIUM

E-mail address: didier.vanderstichelen@uclouvain.be